


MEDICAL PROCEEDINGS

MEDIESE BYDRAES

A South African Journal for the
Advancement of Medical Science

'n Suid-Afrikaanse Tydskrif vir die
Bevordering van die Geneeskunde



Registered at the General Post Office as a Newspaper

By die Hoofposkantoor as Nuisblad Geregistreer

Vol. 2 • No. 8 • 5s

Johannesburg
7 July 1956 7 Julie

Jaarliks £1 : 1 : 0 Yearly

IN THIS ISSUE • IN HIERDIE UITGAWE

Medical Proceedings: A Fortnightly Journal • Electrocardiography Without Tears

Specialist Registration • Cancer Patients

Mediese Bydraes: 'n Tweeweklikse Blad • Elektrokardiografie Sonder Trane

Registrasie van Spesialiteite • Kankerpatiënte

Dysphagia • Hypertension • Neck Swellings • Overbreathing

Preparations and Appliances • Preparate en Toestelle

Notes and News: Berigte • Reviews of Books • Correspondence

Dispensing by Doctors (p. 337)

Index of Contents (p. v)

METIMYD

combined

METICORTELONE • STABILIZED ALBUCID

Schering

CORPORATION



P.O. Box 1010 • Johannesburg: P.O. Box 30 • Cape Town


Uitgewers: Juta en Kie. Bpk.
Posbus 1010 • Johannesburg: Posbus 30 • Kaapstad

INTERCHANGEABLE LUER LOCK SYRINGES

A real **QUALITY PRODUCT** by Everetts **THE** specialists
in the manufacture of Hypodermic equipment.

CLEAR GRADUATIONS

SMOOTH ACTION




REPLACEABLE BARRELS


REPLACEABLE PISTONS

INDESTRUCTIBLE
MARKINGS

PERFECT NEEDLE
Lock



GURR'S "SICO" NEEDLES ARE MADE FOR THESE SYRINGES



Both syringes and needles are made under the same roof ensuring
perfect marriage of needle to syringe.

	1cc.	2cc.	5cc.	10cc.	20cc.	
Prices:—	10/-. —	11/6. —	13/6. 13/6.	17/6. 17/6.	20/- 20/-	Central Nozzles. Side Nozzles.

Stockists and Factory Representatives:

GURR SURGICAL INSTRUMENTS (Pty.) Ltd.

Harley Chambers - Kruis Street
P.O. Box 1562 - Johannesburg

NUUT!

NEW!

METIMYD

STERILE — STERIEL

samestelling

OPHTHALMIC SUSPENSION
OOG-SUSPENSIE

METICORTELONE • ALBUCID Soluble

combined

CONTROLS allergic conditions

CURBS inflammation

CHECKS infection

PROTECTS the eye

5 cc. dropper bottle

METIMYD, brand of prednisolone and sulphacetamide sodium.
METICORTELONE Acetate, brand of prednisolone acetate.
ALBUCID, brand of, sulphacetamide sodium

BEHEER allergiese toestande

BEPERK ontsteking

BEDWING besmetting

BESKERM die oog

5 k.s. indrupbottel

METIMYD, merk van prednisoloon en natrium-sulfasetamied
METICORTELONE-asetaat, merk van prednisoloon-asetaat.
ALBUCID, merk van natrium-sulfasetamied



SCHERAG LTD
JOHANNESBURG

Schering
CORPORATION

**THE NEW SKIN PROTECTIVE
FOR MEDICAL USE
NOW BEING INTRODUCED
IN SOUTH AFRICA IS**

Siopeel

CREAM

It is indicated

- ★ In the management of occupational dermatoses caused by water soluble irritants.
- ★ In post-operative care of the skin following colostomy, ileostomy, haemorrhoidectomy, etc.
- ★ In dealing with the effects of incontinence.
- ★ SIOPEL CREAM incorporates a specially selected silicone. Issued in tubes of 50 grammes and containers of 500 grammes.

IMPERIAL CHEMICAL (PHARMACEUTICALS) LIMITED
WILMSLOW, MANCHESTER (A subsidiary company of Imperial Chemical Industries Limited)

DISTRIBUTED BY:

I.C.I. SOUTH AFRICA (PHARMACEUTICALS) LIMITED
Pan Africa House 77 Troye Street P.O. Box 11270 Johannesburg



Medical Proceedings · Mediese Bydraes

Vol. 2 · No. 8

INDEX · INHOUD

7 July 1956 7 Julie

<i>Editorial: Medical Proceedings—A Fortnightly Journal; Electrocardiography Without Tears; Higher Qualifications for the Registration of Specialists; Treatment and Care of Cancer Patients—Programme of the National Cancer Association of South Africa</i>	317
<i>Redaksioneel: Mediese Bydraes—'n Tweeweklikse Blad; Elektro-kardiografie Sonder Trane; Hoër Kwalifikasies vir die Registrasie van Spesialiteite; Behandeling en Versorging van Kanker pasiënte—Program van die Nasionale Kankervereniging van Suid-Afrika</i>	317
<i>The Modern Treatment of Hypertension: A Review of the New Antihypertensive Drugs. Dr. L. Kornel</i>	319
<i>Dysphagia. Mr. W. L. Phillips, F.R.C.S.</i>	324
<i>Swellings of the Neck: Midline Swellings (Continued from p. 288) Prof. J. H. Louw and Prof. C. F. M. Saint</i>	329

<i>The Hyperventilation or Overbreathing Syndrome. Dr. F. Ames</i> ...	333
<i>Preparations and Appliances: Sigmagen Tablets; Dalton Compound</i> ...	335
<i>Preparate en Toestelle: Sigmagen-Tablette; Dalton-samestelling</i> ...	336
<i>Notes and News: Berigte</i>	337
<i>Reviews of Books: A Text Book of Midwifery for Students and Practitioners (Johnstone); Lectures on the Scientific Basis of Medicine Vol. III, 1953-1954; A Primer of Electrocardiography (Burch and Winsor); Viral and Rickettsial Diseases of the Skin, Eye and Mucous Membranes of Man (Blank and Rake); Yellow Fever Vaccination (Smithburn et al.); Control of Insect Vectors in International Air Traffic: A Survey of Existing Legislation (World Health Organization)</i>	337
<i>Correspondence: Rehabilitation Association for Injured Workmen; Workmen's Rehabilitation Centre (Mr. J. P. Grobbelaar)</i> ...	340



in one tablet

There is no safer potent hypotensor known. No contra-indications and yet effective in $\pm 65\%$ of all cases. Side effects mild and innocuous.



laboratories Africa (Pty) Ltd

4574-2E

ASPIRIN INTOLERANCE IN RHEUMATOID ARTHRITIS

BUFFERIN- THE NEW BETTER TOLERATED ANALGESIC

MOST SUCCESSFUL
ANSWER
YET DISCOVERED

BUFFERIN FAR
BETTER TOLERATED
BY ARTHRITICS

ACTS TWICE AS
QUICKLY AS ASPIRIN

NO EQUIVALENT
IN BRITISH
PHARMACOPOEIA

Following last year's report¹ by the Joint Committee of the Medical Research Council and Nuffield Foundation, many practitioners have returned to aspirin therapy in place of cortisone. Effective therapy with aspirin and similar salicylates unfortunately has been hampered by the problem of gastric irritation, which affects as many as 42% of arthritic patients. Bufferin, the new analgesic which reduces gastric intolerance to a negligible minimum, is the most successful answer to the problem yet discovered.

Bufferin combines acetylsalicylic acid with the antacids aluminium glycinate and magnesium carbonate (gentle buffering acids which do not produce "acid rebound"), and is well tolerated even by arthritics. In blind trials amongst arthritics with proven intolerance to ordinary aspirin, 70% had no gastric symptoms after taking large doses of Bufferin over periods of 4 to 16 months². Previous studies amongst the general population showed that only 1 patient out of 238 had any distress after taking 10 grains of Bufferin.³

Not only is Bufferin better tolerated by the stomach than ordinary aspirin, but its pain relieving ingredient is absorbed twice as quickly into the blood-stream. Trials show that on an average the 10-minute salicylate level after taking Bufferin was more than 20% higher than the 20-minute level after taking ordinary aspirin.³

Bufferin is unique in incorporating aluminium glycinate and magnesium carbonate as buffering agents and has no equivalent in the British Pharmacopoeia.

References: 1. *British Medical Journal*, 1954, 1: 1223. 2. *Journal of the American Medical Association*, June 4, 1955, p. 387. 3. *J. Amer. Pharm. Assoc.* 1950, 39: 21.

Depot Stockists:

Heynes Mathew, Ltd., Cape Town
South African Drug Houses, Durban
E. P. Drug House, Port Elizabeth
James Reid (Pty.) Ltd., Bloemfontein
Cloete Kruger (Pty.) Ltd., P.O. Box
759, Windhoek

Distributed by:

BRISTOL-MYERS (PTY.) LTD.,
P.O. BOX 9706, JOHANNESBURG

BUFFERIN
TRADE MARK
ANTACID ANALGESIC



INTERNATIONAL
medical news bulletin

Current Developments in the Fields of Antibiotics,
Hormones, Nutrition and Clinical Medicine

Prepared for Physicians by the Medical Department of Pfizer International, Inc., 25 Broad Street, New York 4, N.Y., U.S.A.

Vol. III, No. 6, 1956

ANTIBIOTIC NEWS AND NOTES

TERRAMYCIN®* INTRAMUSCULAR "IS BEST CHOICE OF TREATMENT FOR GONORRHEA in penicillin-sensitive patients," according to Goodloe and colleagues.¹ The percentage of "cures" was 74.5 in 59 patients with acute gonorrhea with a dosage of 200 mg. Terramycin Intramuscular, divided into 2 doses injected into each buttock.

ARGENTINA: TERRAMYCIN VAPORIZATIONS RECOMMENDED IN ACUTE RESPIRATORY CONDITIONS - Terramycin vaporizations were administered to 42 children with acute conditions of the respiratory tract. Weinstein² reports good results in 28 patients with acute bronchitis and in 2 with bronchoalveolitis. The treatment "unquestionably relieves" spasmodic bronchitis and is "definitely beneficial to whooping cough." The majority of patients received one daily vaporization of 500 mg. Terramycin diluted in 10 cc. of 75% propylene glycol, by means of an oronasal face mask.

^(R)
TETRACYN**SF† "EXTREMELY EFFECTIVE" IN PEDIATRIC INFECTIONS - An orally administered preparation containing "tetracycline and the vitamin mixture proposed by the National Research Council as a therapeutic adjuvant in conditions of stress" was "extremely effective" in 50 children with a variety of acute pharyngeal and respiratory tract infections, say Andelman and Nathan.³ Treatment was continued until 2-3 days after the patients became afebrile; 5-6 days of therapy were usually sufficient. The authors find it significant that in addition to complete recovery, there were no complicating secondary infections. They feel that "vitamin replacement therapy was of value in hastening the convalescent period" in many cases.

INDIA: "DRAMATIC RESPONSE" TO DIHYDROSTREPTOMYCIN IN RHEUMATIC PERICARDITIS - Banerji⁴ reports a "dramatic response" to dihydrostreptomycin in a patient with rheumatic pericarditis, after penicillin and salicylate treatment had failed. Daily

*Brand of oxytetracycline

**Brand of tetracycline

†Trademark for Pfizer-originated, vitamin-fortified antibiotics

injections of 1 Gm. dihydrostreptomycin, given for 20 days together with multi-vitamins, effected a steadily continued "improvement." In view of the "excellent results" obtained, author recommends that the use of dihydrostreptomycin be explored further in patients with rheumatic carditis refractory to other therapy.

PENICILLIN "SUITABLE" FOR STREPTOCOCCAL MASS PROPHYLAXIS - Studies at the U.S. Naval Training Center in Bainbridge, Md., suggest benzathine penicillin G may be a "suitable agent for mass prophylaxis" against infections due to group A beta-hemolytic streptococci. Brooks and Tilden⁵ found that a single injection (600,000 U.) given to 624 carriers, elicited 597 negative cultures for at least one month thereafter. This according to the authors, "seems to indicate a period of protection of perhaps four to six weeks." They consider it "significant" that rheumatic fever did not occur in any treated recruit.

GERMANY: TETRACYN "EFFECTS A MORE RAPID CURE" IN DYSENTERY than sulfonamides, reports Schütze⁶ in a comparative study between two groups of 40 patients each. In the Tetracyn group 35 were "cured" and 4 "conditionally cured." In the sulfonamide group 28 were "cured" and 3 "conditionally cured." "...Tetracyn effects a more rapid cure than the sulfonamide therapy used up to date...." "In addition to its quicker effectiveness, Tetracyn therapy is preferable to sulfonamides particularly because of its much greater tolerance."

NORWAY: TERRAMYCIN IN RESPIRATORY AND URINARY INFECTIONS - Terramycin given orally produced "good" results in 74 of 96 patients with infections of the respiratory, urinary, gastrointestinal and biliary tracts, the skin and subcutaneous tissues (adenitis, phlebitis), reports Kløvstad.⁷ Dosage was 4 tablets daily, 250 mg. each, given generally for 4-5 days. The patients ranged in age from 10 to over 80 years.

GERMANY: A PENICILLIN-DIHYDROSTREPTOMYCIN COMBINATION* "can cure" various infections caused by a wide range of organisms, such as gonococci, staphylococci, streptococci, pneumococci, enterococci, coli, Friedländer's bacillus and S. pallida. Schubert⁸ observed that 1,000,000 U. of the combination, given 1-3 times, elicited "complete inhibition" of etiologic agents. The antibiotic combination was also used with "good result" in 2 patients with primary and latent syphilis. "...this new combination is indicated not only for short term, but also, especially, for long-range therapy."

PANAMA: TETRACYCLINE IN NONSPECIFIC URETHRITIS - "Cure" was obtained in 27 of 37 patients with nonspecific urethritis after 16 to 48 tablets were administered for 4-12 days, reports Fabrega.⁹ The author thought that "its low toxicity and wide bactericidal spectrum" might make it "an antibiotic par excellence" in the treatment of this disease.

FRANCE: "REMARKABLE" EFFECT OF CHEMOTHERAPY ON RENAL TUBERCULOSIS IN PREGNANCY - Chemotherapy had a "remarkable, even curative, effect on renal tuberculosis..." in 3 pregnant women, observe Aboulker and Muhlrad.¹⁰ The treatment had no bearing on the course of pregnancy and induced no fetopathy. The women received a total of 63 Gm., 35 Gm., and 32 Gm. isoniazid each, after pregnancy periods of respectively 2 months, 4½ months (also 37 Gm. streptomycin and 110 Gm. PAS), and 5½ months (also 10 Gm. dihydrostreptomycin sulfate).

*Available from Pfizer as Combiotic®

BIBLIOGRAPHY

1. Goodloe, O. M., and Roth, L. J.:
Ohio State M. J. 52:34.(Jan.) 1956.
2. Weinstein, M.: Dia méd 27: 2882
(Nov. 24) 1955.
3. Andelman, M. B., and Nathan, L. A.:
Antibiotic Med. 2:45 (Jan.) 1956.
4. Banerji, D.: Calcutta M. J. 52:339
(Nov.) 1955.
5. Brooks, T. J., and Tilden, I. M.:
J.A.M.A. 160:162 (Jan. 21) 1956.
6. Schütze, E.: Therap. Gegenw. 95:7
(Jan.) 1956.
7. Kløvstad, O.: Tidsskr. f. d. norske
laegefor. 76:7 (Jan. 1) 1956.
8. Schubert, E.: Ztschr. f. Haut. u
Geschlechtskr. 20:80 (Feb. 1) 1956.
9. Fabrega, R. D.: Arch. med. paname.
4:161 (July-Aug.-Sept.) 1955.
10. Aboulker, P., and Muhlrad, S.:
Presse méd. 64:45 (Jan. 11) 1956.

HORMONES

DELTACORTIL® "EXCELLENT" IN SEVERE ALLERGY - Johnston and Cazort¹ report the effectiveness of prednisolone in severe allergies. Of 201 patients with severe allergic diseases treated with 232 courses of the steroid, 148 showed "excellent" and 55 "good" results. In addition, "excellent" results were seen in 20 and "good" results in 4 of 25 patients with severe refractory ragweed hay fever.

ITALY: PREDNISOLONE, PREDNISONE REPORTS FROM ITALIAN UNIVERSITIES - Reports from the Universities of Milan, Naples, Pisa, Florence, and Genoa, from hospitals in Brescia and Novara, and from the National Institute for Social Service in Sondalo² unanimously attest to the therapeutic effectiveness and excellent tolerance of prednisolone and prednisone. Prednisone, specifically, was found to be beneficial and effective in mesenchymal neoplasia, malignant neoplasia, lympho-granuloma, leukemia and other hemopathies, Micheli syndrome, nephrotic syndromes, tubercular pleuritis, pemphigus and other dermatoses, histamine cephalalgia, cortisone-resistant ocular diseases, and in gynecologic complaints as well as in the pediatric field.

SWITZERLAND: FIRST SYMPOSIUM ON PREDNISONE IN SWITZERLAND - Clinicians participating in the first prednisone symposium³ confirmed the "indisputable therapeutic value" of the steroid. Böni (U. of Zurich) discusses its "especially strong" anti-inflammatory and antirheumatic effect on 22 patients with progressive chronic polyarthritis. Alphonse (U. of Geneva) reports on the "effectiveness" of prednisone in 5 patients with lupus erythematosus disseminatus; the steroid was used in doses equal to or smaller than the usual amounts of cortisone. In comparing the effects of cortisone and prednisone on epidemic hepatitis, Koller and colleagues (Zurich) conclude that prednisone "is particularly well suited for the treatment of epidemic hepatitis." Daily doses of 30 mg. prednisone "seem to have the same effect as 200 mg. cortisone." A "surprising improvement of the general condition" is seen, appetite returns, and "it is possible to change more rapidly to a protective high-protein diet for the liver."

GERMANY: PREDNISOLONE, PREDNISONE THERAPEUTIC "ADVANCE" IN ADDISON'S DISEASE - Prednisolone and prednisone "... achieve subjectively and objectively a complete clearing of the manifestations of severe Addison's disease," states Overzier.⁴ Three

*Brand of prednisolone in tablets, white, scored, 5mg. tablets, bottles of 10, 20 and 100; in the familiar Pfizer oval shape.

women treated orally with both steroids, as well as with high doses of vitamin C, showed subjective and objective improvement after 3-4 days. The author considers this "...an advance because controlled ambulatory treatment without dietary restrictions is possible for the patient...."

LOCAL HYDROCORTISONE "SUPERIOR" IN PERIARTICULAR PAIN - Hydrocortisone acetate injections into painful joints effected results "superior to those obtained by methods available prior to the advent of hydrocortisone," report Kilroy.⁵ One or more injections (or infiltrations) of 25, 50 or 75 mg. hydrocortisone into the point of greatest tenderness produced "good" results in 36 of 48 patients with tendinitis of the musculocutaneous cuff; complete freedom of pain was felt within 36 to 48 hours. "Good" effects were also obtained in 12 of 21 patients with epicondylitis and in 5 of 6 patients with Quervain's disease. [See note (1) below.]

A report by Grewe⁶ confirms the superiority of hydrocortisone in epicondylitis or styloiditis. Ninety per cent of 33 patients were free of complaints 3-6 months after therapy. Single injections of 25 mg. hydrocortisone acetate were usually sufficient to alleviate the pain. "Excellent results were also seen in the treatment of bursitis."

NOTE (1): Available as Cortril® Acetate Aqueous Suspension containing 25 mg. hydrocortisone acetate per cc.

BIBLIOGRAPHY

- | | |
|--|---|
| 1. Personal communication. | 5. Kilroy, D. O.: California Med. <u>83</u> : |
| 2. Minerva med. <u>46</u> :1693-1760 (Dec. 12) 1955. | 416 (Dec.) 1955. |
| 3. Méd. et hyg. <u>13</u> :465 (Dec. 10) 1955. | 6. Grewe, H. E.: Zentralbl. Chir. <u>80</u> : |
| 4. Overzier, C.: Arztl. Wchnschr. <u>10</u> : 1173 (Dec. 30) 1955. | 1210 (no. 30) 1955. |

TERRAMYCIN* $\frac{S}{F}$ † (brand of oxytetracycline with vitamins)

TETRACYN* $\frac{S}{F}$ † (brand of tetracycline with vitamins)

THERAPY FOR INFECTIOUS DISEASE and THERAPY FOR CONVALESCENCE go hand in hand in formulations, originated by Pfizer, of specific potent antibiotics plus those vitamins necessary to compensate for nutrient loss, strengthen the body's defense mechanisms, and facilitate return to normal.

Available in 250 mg. capsules.

Address for Medical Enquiries:

PFIZER LABORATORIES SOUTH AFRICA (PTY.) LTD.

P.O. Box 7324, Johannesburg.



*Je le pensay,
Dieu le guarit'*

(I TREATED IT, GOD HEALED IT)

It is an odd misnomer that medicine should have become known as The Art of Healing. For it is as true today as it was when Paré coined his famous phrase, that no doctor can heal a wound. All he can do is to eliminate those things that would otherwise delay healing, and avoid the use of medicaments that retard the natural process.


'Furacin' Soluble Ointment satisfies this double need. As effective against bacteria as the antibiotics, it does not delay natural healing or re-growth in any way. Moreover, the *nitrofurans*, the group of chemotherapeutic agents to which 'Furacin' (nitrofurazone) belongs, are unrelated to either the antibiotics or the sulphonamides. There is, therefore, no risk of cross-resistance or cross-sensitization.

FURACIN SOLUBLE OINTMENT

for the treatment and prevention of wound infection
effective . . . convenient . . . economical

In 1 oz. tubes • Samples and literature on request

M. & J. Pharmaceuticals (Pty.) Ltd., Diesel Street, Port Elizabeth




As sure as the dawn

On injecting Cytamen—crystalline vitamin B₁₂—a doctor can be as confident of that injection as he is in the certainty of dawn. Confident that each dose is an exact weight of crystalline vitamin B₁₂ . . . that the stated potency cannot vary . . . that varied response on the part of the patient therefore is not and cannot be due to the product . . . that he has at his command a range of potencies which allows complete flexibility of treatment. In short, Cytamen puts into the doctor's hand a most therapeutically efficient and the most economical means of combating pernicious and other macrocytic anaemias.

CYTAMEN*

Crystalline vitamin B₁₂

50 micrograms 6 x 1 cc., 100 x 1 cc. ampoules and 10 cc. vials
100 micrograms 6 x 1 cc., ampoules and 10 cc. vials
1000 micrograms 3 x 1 cc. ampoules



*When convalescence drags and they
shake their heads at everything, one
naturally thinks of Cytaccon**

Cytaccon Oral Vitamin B₁₂—

Liquid—25 micrograms per fluid drachm. 6-oz. bottles.

Tablets—10 micrograms
in bottles of 50 and 500.
50 micrograms in bottles of 25

TRADE MARK*



GLAXO LABORATORIES (S.A.) (PTY.) LTD., P.O. BOX 21, WADEVILLE, GERMISTON
Agents M. & J. Pharmaceuticals (Pty.) Ltd., P.O. Box 784, Port Elizabeth.

Cortone*

ACETATE (Cortisone Acetate of Merck & Co., Inc.)

...in Oral, Ophthalmic, and Parenteral forms...provides needed dosage and administration flexibility for numerous indications.



TABLETS:
25 mg. in bottles of 20 and 40.



OPHTHALMIC: 0.5% Sterile Suspension of CORTONE Acetate. 2.5% Sterile Suspension of CORTONE Acetate. 1.5% ointment, 3.5-Gm. tube.



PARENTERAL:
20-cc. vial—25 mg./cc.

CORTONE is the original brand of compound E. This product was first made available to the world through Merck & Co., Inc. research and production.

CORTONE ophthalmic preparations provide dramatic benefits in inflammatory ocular disease of the anterior segment; local therapy proves effective, convenient, and economical.

CORTONE in oral and parenteral forms, provides striking control and alleviation of many serious arthritic conditions; also is highly effective in the treatment of certain eye and skin diseases, allergies, and many other disorders.

CORTONE dosage and administration does not neces-

sitate taking any measures not readily available to the physician in everyday practice.

From hundreds of reports in medical publications the world over, comes word of the ever-increasing usefulness of **CORTONE** products. Ever since this compound was made available by Merck & Co., Inc., a growing number of indications have been added to the list of conditions which respond to **CORTONE** in its various forms...more than can be listed here. Literature on the dosage and administration of **CORTONE** for indications encountered in daily practice available on request.

All **CORTONE** tablets carry this trade-mark:

4

Enquiries: P.O. Box 5933, Johannesburg

MERCK-SHARP & DOHME INTERNATIONAL

DIVISION OF MERCK & CO., INC.

161 AVENUE OF THE AMERICAS NEW YORK 13, N.Y., U.S.A.

*CORTONE is the registered trade-mark of Merck & Co., Inc., for its brand of cortisone.



OF ATTACKS OF **ANGINA PECTORIS**

METAMINE tablets (Triethanolamine Trinitrate Biphosphate) *prevent* attacks of angina pectoris or greatly diminish their number and severity.

METAMINE tablets have a duration of action 2-4 times that of Nitroglycerine and are singularly free from side-effects and toxic manifestations.

For full information and literature write to:

KEATINGS PHARMACEUTICALS LIMITED

P.O. BOX 256, JOHANNESBURG

P.O. BOX 568, CAPE TOWN • P.O. BOX 2383, DURBAN • P.O. BOX 789, PORT ELIZABETH

(337)

NEWS!




*Streptokinase-Streptodornase Lederle

Intramuscular VARIDASE^{*} for control of inflammation

MULTIPLE LACERATIONS AND ABRASIONS FOLLOWING ASSAULT



On admission.

24 hours after treatment was initiated;
two Varidase injections have been received.48 hours after admission; four Varidase
injections have been given.

Injected intramuscularly, VARIDASE has produced remarkable results in the treatment of abscess, cellulitis, oedema, epididymitis, hemarthrosis, sinusitis and thrombophlebitis with or without superimposed infection.


VARIDASE quickly dissipates simple inflammation. In treating infected lesions, antibiotics should be administered concomitantly. In such cases VARIDASE breaks down the 'limiting membrane' which contains the infection, allowing passage of the antibiotic.

LEDERLE LABORATORIES DIVISION

AMERICAN CYANAMID COMPANY 30 Rockefeller Plaza New York

Sole Distributors: ALEX LIPWORTH LTD., Johannesburg, Cape Town, Durban & Salisbury

72 hours after admission; treatment
completed.



To assure clinically precise electrocardiograms

... more and more doctors are equipping their offices with the famous Beck-Lee *Cardi-all*

The *Cardi-all* not only provides the physician with heart records of high fidelity and precision but also is so simple to operate, so easy to carry, so free from maintenance cares, so amazingly low in price!

THE NEW PORTABLE BECK-LEE

Cardi-all

DIRECT WRITING ELECTROCARDIOGRAPH

* This is one of a series of advertisements designed to focus attention on the many exclusive features of the *Cardi-all* which the world's largest exclusive manufacturer of electrocardiographs brings to electrocardiography.



Medical Distributors*

SPECIALISTS FOR PHYSICAL MEDICINE APPLIANCES
P.O. Box 3378 JOHANNESBURG Tel. 23-8106

Telegraphic Address: "DISMED"

Office and Showroom at 236, JEPPE STREET

Complete with
all accessories

£297 : 10 : 0

Weight: 27 lb.

A FEW OF THE MANY FEATURES

Clinical Precision—Accurate, dependable cardiograms assured by full frequency Galvo-motor*.

Easy Operation—With less than an hour of training anyone can operate the *Cardi-all* and make a complete record in only a few minutes. Through an ingeniously designed recording unit, change of paper roll can be made in about 10 seconds.

Full Portability—The *Cardi-all* weighs only 27 pounds, completely loaded and all accessories included.

Automatic Control—Specially designed circuits virtually eliminate any stylus wandering due to line voltage variations. An adequately stable iso-electric line for diagnostic purposes is provided by the *Cardi-all*. AC interference done away with by automatic self-grounding circuits plus electronic separation of interfering voltages.

Economically Priced—The *Cardi-all*, complete with all accessories, is only £297:10:0

Literature available from

The Climacteric

"... a mixture of the male and female hormones was more efficacious than oestrogen alone." Brit. Med. Journ. 1953 (July 25), 2,214

The association of methyltestosterone and ethinylœstradiol in Mepilin produces a more complete response in the treatment of menopausal disorders than can be obtained by the use of oestrogens alone.

The presence of methyltestosterone enables a reduction in oestrogen dosage to be made; thus undesirable side effects such as breast turgidity and pelvic congestion are avoided and the risk of withdrawal bleeding is reduced. An increased feeling of confidence and well-being is produced which is both mental and physical.

'MEPILIN' TABLETS & ELIXIR

TRADE MARK

Each teaspoonful (4 ml.) of Mepilin Elixir and each Mepilin Tablet contains ethinylœstradiol 0.01 mg. and methyltestosterone 3 mg.

'MEPILIN' TABLETS

Bottle of 25 at 7/4 and 100 at 23/8

'MEPILIN' ELIXIR

Bottle of 4 fl. oz. at 10/2 and 20 fl. oz. at 38/5

Prices to the Medical Profession

DOSAGE: Menopause and geriatric conditions: average cases — 3 tablets or 3 teaspoonfuls daily. Premenstrual tension and dysmenorrhœa — 2 tablets or 2 teaspoonfuls daily from 10th to 22nd day of the menstrual cycle.

Literature and specimen packings are available on request

BRITISH DRUG HOUSES (SOUTH AFRICA) (PTY.) LTD.

MBP/M/4



*A year of
androgenic effect
in a dozen
injections of*

DEPO*TESTOSTERONE

(Cyclopentylpropionate)

Prolonged: Each injection provides anabolic and androgenic effects up to 4 weeks—longer lasting than any other androgen. **Potent:** This action is maintained at a high level throughout this period. **Economical:** Fewer injections lower the cost to the patient, require less of the physician's time.

VALUABLE in impotence, infantilism, cryptorchidism and premature senility in *men*; menorrhagia, metrorrhagia, painful breast engorgement and control of lactation in *women*; enhances protein anabolism and increases libido in *both sexes*.

Vials of 1 cc. and 10 cc.

Each cc. of Depo-Testosterone contains:

Testosterone cyclopentylpropionate	100 mg.
Chlorbutanol	5 mg.
Cottonseed oil	q.s.

*Trademark

Upjohn

Fine pharmaceuticals since 1886

THE UPJOHN COMPANY, Kalamazoo, Michigan, U.S.A.

Westdene Products (Pty.) Ltd.,

Cape Town: 408 Grand Parade Centre, Castle Street.

Durban: 66/67 National Mutual Buildings, Corner Smith and Gardiner Streets.

Johannesburg: 22-24 Essanby House, 175 Jeppe Street.

Pretoria: 210 Medical Centre, Pretorius Street.

MEDICAL PROCEEDINGS

MEDIESE BYDRAES

A South African Journal for the
Advancement of Medical Science

'n Suid-Afrikaanse Tydskrif vir die
Bevordering van die Geneeskunde

P.O. Box 1010 · Johannesburg | Posbus 1010 · Johannesburg

Vol. 2

July 7 1956 Julie 7

No. 8

EDITORIAL · REDAKSIONEEL

MEDICAL PROCEEDINGS

A FORTNIGHTLY JOURNAL

With this issue *Medical Proceedings* celebrates its first birthday by becoming a fortnightly publication. This has been determined by the truly remarkable support it has received in every respect necessary for successful clinical journalism.

Most of us harbour a Caxtonian curiosity about the techniques of printing. Our readers will, therefore, probably be interested to learn that the cover of the Journal is printed by Verkotype. This produces the decorative, raised type which graces the Journal cover. A solution of many technical, mass production and aesthetic problems made the adoption of this process possible.

The Journal will now also have opportunities to comment more topically on matters of current moment to the profession.

Despite a doubling of its publication frequency, the yearly subscription will remain unaltered at one guinea.

ELECTROCARDIOGRAPHY WITHOUT TEARS

It is clear from the numerous communications we have received that the recent series of articles in this Journal by Dr. L. Schamroth successfully introduced many practitioners to the principles of unipolar electrocardiography by means of one of the simplest and most instructive accounts available.

Dr. Schamroth achieved a unique success in combining simplicity with accuracy. The functional illustrations contributed considerably to the success of the exposition.

The interest in and the demand for Dr. Schamroth's articles have been so great that

MEDIESE BYDRAES

'N TWEEWEEKLIKSE BLAD

Met hierdie uitgawe vier *Mediese Bydraes* sy eerste verjaardag deur 'n tweeweklikse blad te word. Dit is die regstreekse gevolg van die werklik merkwaardige steun wat dit ontvang het in iedersfeer wat vir suksesvolle kliniese joernalistiek noodsaaklik is.

By die meeste van ons bestaan daar 'n Caxtoniaanse nuuskierigheid oor die tegniek van die drukkerskuns. Ons lesers sal derhalwe waarskynlik belang stel in die feit dat die omslag van hierdie Tydskrif in *Verkotype* gedruk is. Dit gee ons die dekoratiewe, verheue lettersoort wat ons voorblad tans versier. 'n Oplossing van talle tegniese, massaproduksie- en estetiese probleme het die toepassing van hierdie tegniek moontlik gemaak.

Die Tydskrif sal nou ook geleentheid kry om meer aktuele kommentaar uit te oefen op aangeent-hede wat die aandag van die professie in beslag neem.

Ondanks die verdubbeling van die publikasie-frekwensie bly die jaarlikse intekengeld onveranderd op een ghienie staan.

ELEKTROKARDIOGRAFIE SONDER TRANE

Uit die talle mededelings wat ons ontvang het, blyk duidelik dat die reeks artikels deur dr. L. Schamroth wat onlangs in hierdie Tydskrif verskyn het, die beginsels van eenpolige elektro-kardiografie op 'n hoogs geslaagde en tog eenvoudige en leersame manier onder die aandag van mediese praktisyns gebring het.

Dr. Schamroth het unieke welslae behaal deur eenvoud met akkuraatheid te verenig. Die funksionele illustrasies het heelwat tot die sukses van die uiteensetting bygedra.

Die belangstelling in en die aanvraag om dr. Schamroth se artikels was so groot dat die

the author has included the material which appeared in this Journal in a monograph which has just been published.*

The principles outlined in the earlier chapters published in this Journal are applied to an understanding of the disorders of cardiac rhythm, as recorded in the electrocardiogram. Self-explanatory diagrams once again characterize the new material in the book and the publisher's claim that no specialized knowledge is needed for an understanding of this account of electrocardiography can be endorsed without reservation.

The volume should be invaluable to undergraduate students. It provides practising doctors with a satisfactory and intelligible introduction to what is generally found to be a complex and difficult subject.

HIGHER QUALIFICATIONS

FOR THE REGISTRATION OF SPECIALISTS

At its meeting in March 1956, in accepting a new list of higher qualifications for the registration of specialties, the Council adopted a new principle. The effect is to entitle a medical practitioner who holds a higher qualification in medicine or in surgery (e.g. the M.D., the M.R.C.P. (Lond.), or the F.R.C.S., Eng. or Edin.) to have this qualification registered as the higher qualification required for registration in one or other of the special branches of medicine, e.g. pathology, dermatology, ophthalmology, orthopaedics, etc. An M.R.C.P. or an F.R.C.S. will, e.g. now entitle a practitioner to be registered as a specialist in ophthalmology provided that he complies with the other requirements, which include a period of training in an approved teaching department of ophthalmology.

In the case of obstetrics and gynaecology, qualifications such as a Fellowship in Surgery will only be accepted if obstetrics and gynaecology were taken as a special subject.

TREATMENT AND CARE OF CANCER PATIENTS

PROGRAMME OF THE NATIONAL CANCER ASSOCIATION OF SOUTH AFRICA

The aims of the National Cancer Association of South Africa include assistance in developing services for the diagnosis and treatment

skrwyer die stof wat in hierdie Tydskrif verskyn het, in 'n monograaf saamgevat het. Dit is so pas gepubliseer.*

Die beginsels, uiteengesit in die vroeëre hoofstukke soos in hierdie Tydskrif gepubliseer, word toegepas op 'n begrip van die versteurings van die hartritme, soos in die elektrocardiogram aangeteken. 'n Kenmerk van die nuwe materiaal in die boek is weer eens selfverduidelikende tekeninge, en die uitgewer se aanspraak dat geen gespesialiseerde kennis nodig is vir 'n begrip van hierdie uiteensetting van elektrocardiografie nie, kan sonder enige voorbehoude beaam word.

Hierdie boekie sal van onskatbare waarde vir ongegradeerdees wees. Dit voorsien praktiserende geneeshere van 'n bevredigende en verstaanbare inleiding tot 'n onderwerp wat allereë as ingewikkeld en moeilik beskou word.

HOËR KWALIFIKASIES

VIR DIE REGISTRASIE VAN SPESIALITEITE

Op sy vergadering in Maart 1956 het die Raad 'n nuwe beginsel neergelê toe hy 'n nuwe lys van hoër kwalifikasies vir die registrasie van spesialiteite aanvaar het. Ten gevolge daarvan sal 'n mediese praktisyn wat in besit is van 'n hoër kwalifikasie in die geneeskunde of in die chirurgie (bv. die M.D., die M.R.C.P. (Lond.) of die F.R.C.S., Eng. of Edin.) geregtig wees om hierdie kwalifikasie te laat registreer as die hoër kwalifikasie wat vereis word vir registrasie in die een of ander spesiale vertakking van die geneeskunde, bv. patologie, dermatologie, oftalmologie, ortopedie, ens. 'n M.R.C.P. of 'n F.R.C.S. sal 'n praktisyn, bv. voortaan geregtig maak om hom as 'n spesialis in oftalmologie te laat registreer mits hy ook aan die ander vereistes voldoen—onder meer 'n opleidingstydperk in 'n goedgekeurde onderrigafdeling vir oftalmologie.

In die geval van verloskunde en ginekologie sal kwalifikasies soos 'n Genootskap in Chirurgie alleen aanvaar word indien die verloskunde en ginekologie as 'n spesiale onderwerp bestudeer is.

BEHANDELING EN VERSORGING VAN KANKERPASIËNTE

PROGRAM VAN DIE NASIONALE KANKER-VERENIGING VAN SUID-AFRIKA

Die oogmerke van die Nasionale Kankervereniging van Suid-Afrika behels ondermeer die verlening van bystand aan publieke hospitale

* *An Introduction to Electrocardiography.* By Dr. L. Schamroth. (1956. Pp. 60 + xi. 106 Illustrations. 21s.). Cape Town: Juta & Company Ltd.

* *An Introduction to Electrocardiography.* Deur dr. L. Schamroth. (1956. 60 bl. + xi. 106 illustrasies. 21s.). Kaapstad: Juta & Kie., Bpk.

of cancer in all public hospitals in the Union. Members of the medical profession are in a key position to help this praiseworthy objective by guiding the Cancer Association in their task.

The Cancer Association is conducting a survey of existing facilities for diagnosis and the various forms of treatment, and has invited the profession to supply relevant data, as well as to recommend ways of dealing with such problems as inoperable cases, special nursing services, financial aid to clients and their families, etc.

Practitioners who answer the questionnaire recently sent out by the National Cancer Association will assist materially in advancing some of the most important aspects of the diagnosis, treatment and management of malignant disease.

in die Unie met die ontwikkeling van dienste vir die diagnose en behandeling van kanker. Lede van die mediese beroep is by uitstek in 'n posisie om hulp te verleen met hierdie prysenswaardige doel deur aan die Kankervereniging leiding in hulle taak te gee.

Die Kankervereniging is besig met 'n opname van bestaande geriewe vir diagnose en die verskillende vorms van behandeling en het die mediese beroep uitgenooi om toepaslike gegewens te verstrek, sowel as om maniere aan te beveel waarop probleme soos karsinoomgevalle wat nie geopereer kan word nie, spesiale verpleegdienste, finansiële bystand vir pasiënte en hulle gesinne ens., aangepak kan word.

Praktisyns wat die vraelys wat onlangs deur die Kankervereniging uitgestuur is, beantwoord, sal op tasbare wyse bydra tot die bevordering van sommige van die belangrikste aspekte van die diagnose, behandeling en beheer van kanker.

THE MODERN TREATMENT OF HYPERTENSION

A REVIEW OF THE NEW ANTIHYPERTENSIVE DRUGS

L. KORNEL, M.D.*

Medical Outpatient Department 'C', Rothschild Hadassah University Hospital, Jerusalem, Israel

Hypertension with all its more or less dangerous complications is one of the most common diseases of our century. The rapid development and progress of technique connected with the enormous rise in the tempo of life and the intensity of work, permanent tension in office and home and actual lack of psychic relaxation, are all apparently the most important factors responsible for having made hypertensive disease a true plague of modern society. Many efforts have been made during the last few years to find an effective drug against all forms of hypertension.

In view of the multiple causes of hypertension no single drug is likely to be the solution to the therapeutic problem. Psychoneurogenic, cardiovascular, endocrine and renal factors are involved, often apparently on an hereditary basis. Therefore, in the treatment of hypertension the influence of the environment, the diet and the mind must be considered. In spite of all efforts to influence these factors, very little can be done to stop the development of hypertensive disease or to prevent dangerous complications. The new

hypotensive drugs which have lately appeared seem to introduce a new era in the treatment of hypertension and probably mean the beginning of a real success in the fight against hypertensive diseases.

Drugs of this type which call for special attention are: Hexamethonium, Veriloid, Apresoline and Rauwolfia. We intend to describe each of them briefly: their pharmacology, mode of action, clinical use with results obtained and chiefly their application in the case of outpatients continuing their daily activities.

HEXAMETHONIUM AND SIMILAR COMPOUNDS

The production of neurogenic hypertension in experimental animals by surgical procedures (Grimson 1950) suggesting the probability of a nervous factor in the etiology of human hypertension, has focused attention on the use of adrenergic blocking agents in the treatment of this disease. Since the results obtained with the first adrenergic blocking drugs (Priscoline, T.E.A.B.) were not satisfactory, the next step was made with the introduction of hexamethonium bromide by Paton and Zaimis (1949, 1951) as a substance which

* British Council Research Scholar, at present at the Department of Medicine, Queen Elizabeth Hospital, Birmingham, England.

blocks the transmission of nervous impulses across the synapses of autonomic ganglia.

Its structural formula is shown in Fig. 1.




Fig. 1. Hexamethylene-1, 6-bis(trimethylammonium).

Its action is not only sympathicolytic but also vagolytic, the latter being one of the disadvantages of this compound.

Many reports and publications have appeared about the clinical use and effectiveness of hexamethonium in the treatment of hypertension. It seems to be an established fact that its benefit in hospitalized patients using the drug intravenously or subcutaneously is quite certain (repeated doses of 25-50 mg.) and the results obtained in the control of hypertension are impressive, but the oral management gives rather disappointing results. The oral dose must be much greater than that given by injections (125-500 mg. per dose repeated every 6-7 hours) and the results vary from day to day, probably due to uneven and inadequate absorption from the gastrointestinal tract.

'Control of blood pressure with oral hexamethonium is unsatisfactory in most cases and oral treatment might well be reserved for those cases with severe headaches but without objective evidence of vascular degeneration. . . . Morrison and also Smirk and Alstad have shown that adequate control of blood pressure even in severe cases can be obtained with parenteral hexamethonium, but the difficulties in controlling this method of treatment with a potentially dangerous drug are such as to limit its general use' (*British Medical Journal*, 1953, 1, 1319). Great care should be taken when complications exist—impaired renal function, coronary artery disease and existing or threatened cerebral vascular accidents.

The commonest of the undesirable effects is postural hypotension with all its accompanying symptoms; also blurred vision, due to pupillary dilatation, unpleasant sensations like dryness of the mouth, nausea, vomiting, drowsiness and constipation. Paralytic ileus may develop, but rarely, and complaints of transient impotence or loss of ejaculatory power are not uncommon. Tolerance to the drug may develop rapidly, making large doses for continuous treatment necessary.

All these circumstances have stimulated the search for other ganglion-blocking drugs which would obviate the difficulties and disadvantages of hexamethonium, especially in oral treatment. Recently, a few articles have appeared attempting to estimate and compare the effectiveness of new sympathicolytic agents of the methonium series. Particular attention should be paid to phenyldimethonium and pentapyrrolidinium, which seem to overcome some of the disadvantages of hexamethonium. Further research is required.

VERILOID AND OTHER VERATRUM PREPARATIONS

The plants *Veratrum viride* and *Veratrum album* were already known 80 years ago for their hypotensive action, but because of their severe toxicity they could not be used in practice. In the last few years many attempts have been made to obtain extracts of these plants free from their toxic side effects.

Veriloid* is the purified, biologically standardized extract of alkaloids of *Veratrum viride*. This extract is itself a mixture of several alkaloids amongst which the most active are protoveratrine, germerine and cryptenamine. Its effect is the result of its action on at least 2 different sites in the body:

1. A stimulatory effect on afferent vagal nerve endings in the wall of the left ventricle with reflex action on the efferent vagal fibres resulting in bradycardia. This bradycardia may be inhibited by atropine or vagotomy (von Bezold reflex, 1867).

2. A direct central action independent of vagal stimulation, shown by perfusion of the isolated head in cross-circulation experiments in the dog (Stutzman, 1951). This action is unaffected by atropine or bilateral vagotomy. Apart from this, certain of the veratrum alkaloids have a cardiotonic effect on the failing mammalian heart, like the cardiac glycosides, increasing its efficiency, work performance and oxygen consumption.

Veriloid has been used recently in the treatment of hypertensive diseases and many reports and publications have appeared during the last 3 years indicating its high effectiveness in the control of hypertension, especially its malignant form. Good results have been obtained by parenteral as well as by oral administration. The latter method permits wide application of Veriloid in the treatment of outpatients continuing their daily activities and work. Unfortunately this drug, too, may

* Produced by the Riker Laboratories.

cause toxic symptoms and great care must be taken in the adjustment of the dosage, reserving a strong individual approach for each patient. The excessive fall of blood pressure may even lead to collapse. Ephedrine *per os* or subcutaneously was found to be an effective antidote and this fact considerably diminishes the aforementioned dangerous disadvantage. The main toxic effects are a feeling of weakness and drowsiness, nausea and vomiting. The limits between the toxic and the effective but non-toxic doses are very narrow and this calls for precise and cautious dosage adjustment.

Our personal experience showed that in a very high percentage of cases the appropriate dosage can be obtained by:

1. Cautious adjustment of an optimal single dose (intake of the drug followed by $\frac{1}{2}$ -hourly or hourly measurement of the blood pressure, allowing the peak of action to be determined and consequently the time of absorption and the excretion of Veriloid);

2. Optimal spacing of further doses during the day. This very exact titration easily helps to find the dose between these limits mentioned above. Moreover, we are convinced that some of the reported disappointing results are due to inexact dosage adjustment. The average interval between 2 orally administered doses is 5-6 hours and the daily dose varies considerably from 6-20 mg. and even more. A significant fall in both the systolic and the diastolic pressures is noticeable and the subjective and objective symptoms improve quickly. Tolerance to the drug has not been noted even during quite prolonged treatment.

The parenteral use of Veriloid finds its proper application in hospitalized patients, especially in pre-eclampsia and eclampsia gravidarum. The same purified extract of a mixture of alkaloids of *Veratrum viride* is represented by Veradine.*

Among the other veratrum preparations should be mentioned the pure crystalline simple alkaloid of veratrum-protoveratrine. This is a very potent hypotensive agent, but owing to its very low emetic : therapeutic ratio (1:1) it cannot find wide application. Cryptenamine† has been announced as the only isolated veratrum alkaloid preparation with 4 times the range of therapeutic safety, but so far very little has been reported about it.

* The product of Teva Middle East Pharmaceutical and Chemical Works Ltd., Jerusalem, Israel.

† Unitensin of Irvin Neisler Co., U.S.A.

APRESOLINE

This drug is the derivative of phthalazine. Its structural formula is shown in Fig. 2.




Fig. 2. 1-hydrazinophthalazine.

One of the first who studied its pharmacological properties and actions was Schroeder who was able to block the pressor action of pherentasin with this compound. According to his reports and those of other investigators (Freis, Gross, Meier, Moyer and others) the hypotensive response to Apresoline is due to 3 mechanisms:

1. *The central action.* Experimental evidence indicates its chief locus to be in the midbrain, preventing an excessive outflow of sympathetic vasopressor impulses.

2. *Moderate adrenergic blockade* by inhibiting the pressor action of nor-epinephrine and epinephrine.

3. *Antagonistic action against hypertensive substances* considered to be of basic importance in various forms of hypertension: angiotonin (hypertensin), serotonin and pherentasin.

The most significant pharmacological actions of Apresoline are the increase in renal blood flow and the inhibition of cerebral hormonal vasopressor substances. Both actions are highly desirable in the therapy of hypertensive disease. No other substance is known to have such an effect on the cerebral pressor substances, and this is believed to explain the effectiveness of hydrazinophthalazine in patients who have neurogenic hypertension and have failed to benefit from extensive lumbodorsal sympathectomy.

The increase in renal blood flow is due to a decrease in resistance in the renal vascular bed. This, however, does not increase renal function and there is no reason to expect that therapy with this drug will reverse renal failure.

The toxicity of Apresoline has been reported to be quite low. The chief side effects are tachycardia and headaches which may occur at the beginning, but usually disappear in the course of the treatment. Further,

nausea and vomiting, which are uncommon toxic manifestations, may occur, in which case the drug must often be discontinued. The only effective means for overcoming this intolerance is to start with very small doses and increase the amount slowly. Overdosage with Apresoline represents no problem in practice. Both in Man and in experimental animals there is a 'floor' below which blood pressure will not drop despite increasing doses.

Apresoline is available for oral or parenteral use, but its chief and only practical application in prolonged treatment is oral. The dosage schedule should be adjusted to the individual response. It may take 8 weeks or more to obtain optimal benefit. The usual initial dose is 10 or 25 mg. which has to be repeated 4-6 times daily. This dosage should be increased every week till the optimal hypotensive response occurs or the appearance of side effects permits no further dosage increase.

Some patients, after more or less protracted administration, begin to show a tendency to tolerate the drug. This tolerance may be lessened by a due spacing of doses and disappears rather rapidly upon temporary withdrawal of the drug.

l-Hydrazinophthalazine is most effective in hypertension persisting or recurring after sympathectomy. The drug gives also favourable results in benign uncomplicated and early malignant hypertension. When renal damage is advanced, as in chronic renal hypertension and chronic glomerulo-nephritis, the value of Apresoline is considerably diminished.

RAUWOLFIA SERPENTINA

The extracts of the root of the plant *Rauwolfia serpentina*, found chiefly in India and Malaya, have been used in these countries for centuries as a general sedative, as an antidote for the poison of insects and reptiles and as a stimulant to uterine contractions.

Recently this plant has attracted the attention of research workers because of its hypotensive action.

The first purified extracts of total alkaloids of *Rauwolfia* appeared under the name Rauviloid* and Raudixin;† then the most potent separate alkaloids were isolated (Serpasil).‡ Now several preparations under various names are obtainable.

Early research on the root showed that it contained 2 groups of alkaloids—the ajmaline

group and the serpentine group. Its action is chiefly central (on the vasomotor centre), causing generalized vasodilatation resulting in a lowering of blood pressure. General sedation which appears and the feeling of well-being are thought to be very desirable effects.

The hypotensive action of this drug is of moderate intensity and it seems to be the medication of choice in mild hypertension, especially effective in young, labile (neurotic) hypertensive patients without advanced vascular changes. This preparation does not produce undesirable side effects even when given in excessive amounts. Only in a very low percentage of cases were a few mild toxic symptoms described: excessive drowsiness, lassitude, anorexia and nausea, which cease quickly after withdrawal of the drug.

Rauwolfia is given orally. Although the dosage must be individualized, most patients require 4-8 mg. of purified total alkaloids daily, divided into 2-4 doses. The drug has to be taken after meals. The maximum hypotensive effect is generally reached within a week after the beginning of treatment.

When given in conjunction with Veratrum or Pentapyrrolidinium in severe hypertension, the concurrent administration of the 2 drugs effects a response which may well be greater than the sum of the effects of the 2 drugs if given separately. This may indicate the synergistic potentiation which permits the application of relatively smaller doses of both.

Unfortunately, none of the foregoing drugs can be considered the perfect remedy for hypertensive disease, and it should be stressed that no satisfactory studies have yet been made on the combined use of the different drugs, although the few reports which have appeared on Apresoline-Hexamethonium and Rauviloid-Veriloid combinations promise rather encouraging results.

ADDENDUM

Since the preparation of this paper a few reports have appeared, drawing attention to some recently discovered properties of the *l*-hydrazinophthalazine and the *Rauwolfia* drugs, which could limit their use in certain circumstances:

1. Suggestive evidence is presented by Aitchison *et al.* (Brit. Heart J., 1955, 17, 425) that Apresoline causes elevation of the pulmonary arterial pressure and an increase in the cardiac output (simultaneously with the fall in the systemic pressure) in patients with mitral disease. Consequently, when used in cases of hypertension associated with mitral

* Riker Laboratories, Inc.

† Squibb.

‡ Ciba Ltd.

disease, it may aggravate the symptoms of cardiac failure, increasing the load on the right ventricle, or even cause pulmonary oedema.

2. The possible anti-diuretic effect of the Rauwolfia alkaloids, finding its expression in the rapid gain in weight of patients while on treatment, has been studied by McGregor and Segal (Brit. Heart J., 1955, 17, 391). Congestive heart failure may be precipitated in cases with previous myocardial damage, due to fluid retention. Further observations are required.

OPSOMMING

Die veelvoudige oorsake van hipertensie maak dit onwaarskynlik dat enige enkele middel die terapeutiese probleem sal oplos. By die behandeling van hipertensie moet die invloed van omgewing, dieet en die geestesgesteldheid van die pasiënt in ooreenstemming geneem word.

Die nuwe hipotensiemiddels wat in die jongste tyd beskikbaar gestel is, het 'n nuwe tydperk ingelui vir sover dit die behandeling van hipertensie betref.

Die middels van hierdie aard wat spesiale aandag verdien, is: Heksametonium, Veriloid, Apresoline en Rauwolfia.

Die farmakologie, die werkingsmanier, die kliniese gebruik van en die resultate wat verkry is met hierdie middels word in oënskynlike mate, veral wat betref die behandeling van buitepasiënte wat hul gewone daaglikse werksaamhede voortsit.

Die oorsig word afgesluit met 'n besonder volledige bibliografie.

REFERENCES

HEXAMETHONIUM

- Morrison, B. (1953): Brit. Med. J., 1, 1291.
- Morrison, B. and Paton, W. D. M. (1953): *Ibid.*, 1299.
- Paton, W. D. M. and Zaimis, E. J. (1948): Nature, Lond., 162, 810.
- Paton, W. D. M. and Zaimis, E. J. (1949): Brit. J. Pharmacol., 4, 381.
- Paton, W. D. M. and Zaimis, E. J. (1951): *Ibid.*, 6, 155.
- Organe, G., Paton, W. D. M. and Zaimis, E. J. (1949): Lancet, 1, 21.
- Zaimis, E. J. (1950): Brit. J. Pharmacol., 5, 424.
- Moyer, I. H. and Mills, L. C. (1953): J. Clin. Invest., 32, 172.
- Smirk, F. H. and Alstad, K. S. (1951): Brit. Med. J., 1, 1217.
- Schroeder, H. A. (1951): Amer. J. Med., 10, 189.
- Saville, S. (1950): Lancet, 2, 358.
- Freis, E. D., Finnerty, F. A., Schnaper, H. W. and Johnson, R. L. (1952): Circulation (N.Y.), 5, 20.
- Locket, S., Swann, P. G. and Grieve, W. S. M. (1951): Brit. Med. J., 1, 778.
- Locket, S., Swann, P. G., Grieve, W. S. M. and Player, H. P. (1952): *Ibid.*, 1, 254.
- Campbell, A. J. M., Graham, J. G. and Maxwell, R. D. H. (1952): Brit. Med. J., 1, 251.
- Campbell, A. J. M. and Robertson, E. (1950): *Ibid.*, 2, 804.
- Friedman, C. E. (1951): Acta Med. Scand., 140, Suppl. 257.
- Keith, N. M., Nagener, H. A. and Barker, N. W. (1939): Amer. J. Med. Sci., 197, 332.
- Brit. Med. J. (1953): 1, 1319.
- Conway, J. (1953): Brit. Med. J., 1, 814.
- Gilmore, H. L., Kopelman, H., McMichael, J. and Milne, I. G. (1952): Lancet, 2, 898.
- Lyons, R. H., Hoobler, S. W., Neligh, R. B., Moe, G. K. and Peet, M. M. (1948): J. Amer. Med. Assoc., 136, 608.
- Page, I. H. and McCubbin, J. (1951): Circulation, 4, 70.
- Perera, G. A. and Krane, S. M. (1951): Ann. Int. Med., 34, 1017.
- Morris, N. (1953): Lancet, 1, 322.
- Mason, A. A. and Pelmore, J. F. (1953): Brit. Med. J., 1, 250.
- Rosenheim, M. L. (1954): Brit. Med. J., 2, 1181.

OTHER COMPOUNDS OF METHONIUM SERIES

- Wien, R. and Mason, D. F. J. (1953): Lancet, 1, 454.
- Maxwell, R. D. H. and Campbell, A. J. M. (1953): *Ibid.*, 455.
- Smirk, F. H. (1953): *Ibid.*, 457.
- Smirk, F. H. (1955): Ann. Rev. Med., V, 6.

VERATRUM

- Von Bezold, A. and Hirt, L. (1867): Untersuch. Physiol. Lab. Würzburg, 1, 73.
- Bryant, R. D. (1935): Amer. J. Obstet. Gynec., 30, 46.
- Krayer, O., Wood, E. M. and Montes, D. G. (1943): J. Pharmacol. Exper. Therap., 79, 215.
- Meilman, E. and Krayer, O. (1949): J. Clin. Invest., 28, 798.
- Assali, N. S., Moe, G. K. and Mendez, R. (1944): J. Pharmacol. Exp. Therap., 82, 167.
- Assali, N. S. and Pryotowsky, H. (1950): *Ibid.*, 100, 251.
- Dawes, G. S. (1947): *Ibid.*, 89, 325.
- Freis, E. D. and Stanton, I. R. (1948): Amer. Heart J., 36, 723.
- Freis, E. D., Stanton, I. R., Culbertson, I. W., Litter, I., Halperin, M. H., Burnett, C. M. and Wilkins, R. W. (1949): J. Clin. Invest., 28, 353.
- Wilkins, R. W., Stanton, I. R. and Freis, E. D. (1949): Proc. Soc. Exp. Biol. Med., 72, 302.
- Wilkins, R. W., Stanton, I. R., Freis, E. D. and Meisler, F. C. (1950): J. Pharmacol., 98, 166.
- Stutzman, I. W., Maison, G. L. and Kusserour, G. W. (1941): Proc. Soc. Exp. Biol. Med., 71, 725.
- Stutzman, I. W. and Maison, G. L. (1950): Fed. Proc., 9, 318.
- Stutzman, I. W., Simon, H. and Maison, G. L. (1951): J. Pharmacol., 101, 310.
- Swiss, E. D., Bain, R., Maison, G. L. and Stutzman, I. W. (1951): *Ibid.*, 35.
- Swiss, E. D., Simon, H. and Stutzman, I. W. (1951): Feder. Proc., 10, 339.
- Stearns, N. S. and Maison, G. L. (1950): J. Pharmacol., 100, 238.
- Stearns, N. S. and Ellis, L. B. (1952): New Eng. J. Med., 246, 397.
- Kauntze, R. and Trounce, I. (1951): Lancet, 1, 549; 2, 1002.
- Kauntze, R. (1952): Proc. Roy. Soc. Med., 45, 277.
- McNair, I. D., Griffith, G. C. and Elek, S. R. (1950): Amer. Heart J., 40, 382.
- Coe, W. S., Best, M. M. and Kinsman, J. M. (1950): J. Amer. Med. Assoc., 143, 5.
- Josephs, I. L. (1950): Ann. West. Med. Surg., 4, 789.

24. Barrow, J. G. and Sikes, C. R. (1951): *Amer. Heart J.*, **41**, 742.
25. Mills, L. C. and Moyer, I. H. (1952): *A.M.A. Arch. Int. Med.*, **90**, 587.
26. Mills, L. C. and Johnson, L. C. (1952): *Feder. Proc.*, **11**, 360.
27. Moyer, I. H., Miller, S. I., Tashnek, A. B., Snyder, H. and Bowman, R. O. (1953): *Amer. J. Med.*, **14**, 175.
28. Goldman, R. and Frierson, H. R. (1953): *Ibid.*, 168.
29. Davis, E. (1953): *A.M.A. Arch. Int. Med.*, **91**, 715.
30. Kornel, L. and Sklan, B. (1953): *Acta Med. Orient.*, **12**, 326.
12. Page, I. H. (1951): *J. Amer. Med. Assoc.*, **147**, 1311.
13. Schroeder, H. A. (1951): *Amer. J. Med.*, **10**, 189.
14. Schroeder, H. A. (1952): *Circulation*, **5**, 28.
15. Schroeder, H. A. (1952): *Arch. Int. Med.*, **89**, 523.
16. Shapiro and Grollman (1953): *Circulation*, **8**, 188.
17. Taylor, R. D., Page, I. H. and Corcoran, A. C. (1951): *Arch. Int. Med.*, **88**, 1.
18. Riven, S. S., Pocock, B. G., Kory, R. C., Roehm, D. C., Anderson, R. S. and Meneely, G. R. (1953): *Amer. J. Med.*, **14**, 160.
19. Aitchison, J. D., Cranston, W. I. and Priest, E. A. (1955): *Brit. Heart J.*, **17**, 425.

APRESOLINE

1. Barrett, W., Reitze, W., Plummer, A. J. and Yonkman, F. F. (1952): *Fed. Proc.*, April.
2. Britton, J. B., Taylor, I. P. and Ahlquist, R. P. (1951): *Fed. Proc.*, **10**, 282.
3. Craver, B. N., Barrett, W., Cameron, A. and Yonkman, F. F. (1951): *J. Amer. Pharm. A. (Sc. Ed.)*, **40**, 559.
4. Editorial (1951): *J. Amer. Med. Assoc.*, **146**, 1326.
5. Freis, E. D. and Finnerty, F. A. (1950): *Proc. Soc. Exp. Biol. Med.*, **75**, 23.
6. Freis, E. D., McKay, J. C. and Oliver, W. F. (1951): *Circulation*, **3**, 254.
7. Freis, E. D., Rose, Higgins, Finnerty, Kelley and Partenape (1953): *Circulation*, **8**, 199.
8. Grimson, K. S. and Chittum, J. R. (1949): *Amer. J. Med.*, **7**, 251.
9. Grimson, K. S., Chittum, J. R. and Metcalf, B. H. (1950): *Fed. Proc.*, **9**, 279.
10. (1950): *Experientia*, **6**, 19.
11. Moyer, J. H., Handley, C. A. and Higgins, R. A. (1951): *J. Pharmacol. Exper. Therap.*, **103**, 368.

RAUWOLFIA

1. Wilkins, R. W. (1952): *Ann. Intern. Med.*, **37**, 1144.
2. Wilkins, R. W. and Judson, W. E. (1953): *New Engl. J. Med.*, **248**, 48.
3. Wilkins, R. W., Judson, W. E. and Staton, J. R. (1952): *Proc. New Eng. Cardiovasc. Soc.*, 1951-1952, p. 34.
4. Vakil, R. J. (1949): *Brit. Heart J.*, **11**, 350.
5. Meilman, E. (1953): *New Eng. J. Med.*, **248**, 894, 936.
6. Ford, R. V., Livesay, W. R., Miller, S. I. and Mayer, J. H. (1953): *Med. Rec. Ann.*, **47**, 608.
7. Arnold, O. H. and Bock, K. D. (1953): *Dtsch. Med. Wschr.*, **78**, 565.
8. Vida, F. (1952): *Die Med.*, **37**, 1157.
9. Arnold, O. H. (1952): *Ther. d. Gegenw.*, **91**, 167.
10. Seliger, H. (1952): *Ther. d. Gegenw.*, **91**, 411.
11. Chopra, R. N. (1953): *Indigenous Drugs of India*. Calcutta.
12. McGregor, M. and Segal, N. (1955): *Brit. Heart J.*, **17**, 391.

DYSPHAGIA

W. L. PHILLIPS, M.R.C.P., Lond.; F.R.C.S., Eng.

Cape Town

Dysphagia or difficulty in swallowing may result from organic or functional conditions arising anywhere along the pathway from the mouth to the stomach.

Swallowing is initiated when chewed food leaves the mouth and is terminated when it enters the stomach.

THE SWALLOWING MECHANISM

Foods which have been chewed, and fluids, are guided to the back of the mouth by the tongue and the facial muscles, thence to the pharynx. The naso-pharynx is closed off by the palatal muscles so that food will not regurgitate through the nose, and the glottis is occluded to prevent the food and fluid entering the trachea. Finally the victuals are forced into the oesophagus by the constrictor muscles.

It should be understood clearly that the oesophagus is not merely an inert channel

through which food passes, but is in fact an actively constricting and propelling muscular organ. Carefully regulated peristalsis occurs along the entire 10 inches of its adult length up to the point where it finally passes through the diaphragmatic hiatus into the fundus of the stomach.

Inco-ordination of this neuro-muscular mechanism may thus interfere with the swallowing process at any site from the pharyngeal region to the oesophago-gastric junction. If peristalsis is lost, the oesophagus may merely become a channel through which the food drops. If lower sphincteric action is lost, the stomach contents may regurgitate into the oesophagus by anti-peristaltic waves.

The oesophagus is in close relationship to many structures along its course, and abnormalities affecting any of these may interfere with the act of swallowing. The oesophagus

may be pulled or pushed out of place or it may be attacked by any of the lesions which commonly affect hollow muscular viscera.

ETIOLOGY OF DYSPHAGIA

A. LESIONS OF THE PHARYNX

1. *Inflammatory Changes.* Difficulty in swallowing is most frequently caused by an inflammatory lesion such as tonsillitis or pharyngitis. The throat is not only sore, but swallowing itself may be a painful effort. Despite the pain, there is no hold-up of the food.

2. *Paralysis of Muscles.* The muscles of the pharynx must work in a co-ordinated manner in order to receive the bolus of food, close off the naso-pharynx and the larynx and subsequently force the food through the upper opening of the oesophagus. Diphtheritic paralysis of the muscles of the soft palate will interfere with this action. Food, particularly fluids, will then frequently regurgitate through the nose when swallowing is attempted.

Bulbar paralysis complicating poliomyelitis results in an even more extensive muscular weakness, with complete failure of the swallowing mechanism. Fluids enter the glottis and pass into the trachea. Invariably other muscle paralyses are present so that not only is swallowing impossible, but there is an associated loss of the cough reflex.

Myasthenia gravis may be associated with dysphagia, due to actual muscle weakness.

3. *Anaemia.* The dysphagia due to anaemia (as seen in the Plummer-Vinson syndrome) follows on the muscular inco-ordination which affects the upper part of the oesophagus and pharynx.

4. *Pharyngeal Diverticulum.* This results from a weakness between the transverse and oblique fibres of the inferior constrictor muscle of the pharynx. The mucous membrane bulges posteriorly and a sac is formed, which eventually enlarges with a pocketing of food within. This is a hernia and is of quite frequent occurrence (Fig. 1).

5. *Hysteria.* Hysteria may simulate any of these dysphagic patterns. The patient states that swallowing is impossible and often complains of an obstructing lump in the throat—the so-called globus hystericus.

B. LESIONS OF THE UPPER OESOPHAGUS

1. *Foreign Bodies.* The posterior wall of the upper part of the oesophagus seems to be almost the site of election for the impaction



Fig. 1. Three views of a pharyngeal pouch. These show the posterior position and the overhanging pouch causing some compression of the oesophagus.

Fig. 2. An upper denture impacted in the upper oesophagus. The patient was admitted in a very toxic state with severe pain, high temperature and subcutaneous emphysema of the neck.

of fish bones. Dysphagia occurs immediately and dramatically. The individual is usually aware that he has swallowed a bone and every subsequent swallowing movement is painful. It is well known that an abrasion or tear of the mucous membrane of the oesophagus originally produced by the foreign body may lead the patient to insist that the bone is still impacted in his throat (Fig. 2).

2. *Pressure from Outside.* Compression and constriction of the neck structures by an enlarging thyroid gland may produce a difficulty in swallowing of gradual onset. Occasionally a thyroid adenoma becomes impacted in the superior mediastinum and the dysphagia commences abruptly. This presupposes considerable pressure and therefore it is not surprising to find a concomitant dyspnoea. The thyroid may not appear enlarged and it may

be very difficult to detect the presence of a plunging goitre which has become obstructed behind the upper part of the manubrium sterni. Similarly, an aortic aneurysm or the osteophytic outgrowths associated with cervical vertebral osteo-arthritis may cause dysphagia through pressure on the upper end of the oesophagus.

3. *Corrosives.* The swallowing of corrosives (e.g. caustic soda or even strong acids) affects not only the mouth, but often the upper part of the oesophagus. The mouth and lips may evidence corrosive effects. If a large amount of the poison has been swallowed the entire length of the oesophagus may be burnt.

4. *Carcinoma.* Carcinoma of the upper part of the oesophagus usually obstructs the lumen of the oesophagus gradually, though a sudden complete occlusion may be caused by blockage due to a small piece of food. It is not the commonest site for the development of a carcinoma which, when it does attack this region, is of slow development, leading to impaired swallowing and finally, in the absence of treatment, only allowing the passage of fluids.

C. LESIONS OF THE MID-OESOPHAGUS

1. *Carcinoma.* The most frequently occurring mid-oesophageal lesion causing dysphagia is a carcinoma which may arise in the

oesophagus itself or be an extension from the left main bronchus. The lesion usually occurs at the site of closest relationship between the aorta and oesophagus, i.e. where the arch of the aorta crosses the oesophagus.

2. *Pressure from Aneurysm.* The oesophagus is crossed by the aortic arch; thus any enlargement of this main vessel will obstruct it. Aneurysm of the arch of the aorta leads to symptoms not only attributable to the oesophageal obstruction, but also to paralysis of the recurrent laryngeal nerve as it hooks around the vessel. The upper mediastinum is a relatively small region; consequently the encroachment by the aneurysm soon leads to compression of the adjacent structures (Fig. 3).

Aberrant vascular rings may cause dysphagia, occurring in the early months of life.

3. *Pressure from Mediastinal Tumours.* Any of the various mediastinal tumours may cause compression in a manner similar to the foregoing. It is surprising to discover how often a tumour may be lodged in the mediastinum without causing any dysphagic symptoms. The rate of enlargement is obviously the important factor, because a slowly growing tumour must suffer considerable enlargement before causing obstruction of the oesophagus. The patient may complain of some slight difficulty in swallowing, but more often than not the symp-




Fig. 3. A large aneurysm of the ascending aorta causing partial obstruction of the oesophagus.

Fig. 4. A superior mediastinal thyroid cyst causing displacement of the oesophagus with dysphagia.

tom is ignored until the size of the tumour causes obvious mechanical obstructive effects (Fig. 4).

4. *Mediastinitis*. Infection of the mediastinum is most often secondary to a penetrating lesion of the oesophagus or bronchus. The cause of the penetrating injury may be quite apparent and initially dysphagia may be absent. With the development of a mediastinitis a difficulty in swallowing ensues, because the muscular tube becomes surrounded by the active inflammatory process.

5. *Empyema*. Similarly, a large empyema may cause swallowing difficulties. If the pus is under tension, the dysphagia is due to direct compression of the oesophagus. In more chronic cases the thickening of the pleura and the formation of a pyogenic membrane provides a firm rigid lining on one side of the oesophagus. The patient can swallow, but with considerable embarrassment.

D. LESIONS OF THE LOWER END OF THE OESOPHAGUS

1. *Carcinoma*. Carcinoma on the lower end of the oesophagus is the commonest cause of dysphagia in the age group 45–60 years. The usual histology is that of a squamous-celled

carcinoma, which may start as a submucous lesion. If, however, the carcinoma arises in a gland, it will have an adenomatous structure large enough to obstruct the lumen; or a tumour surrounding the oesophagus may cause a narrowing or constricting effect.

Occasionally a carcinoma of the cardia of the stomach extends upwards, leading to obstruction of the lower part of the oesophagus.

2. *Achalasia*. Achalasia or cardiospasm has been attributed to several causes, of which neuro-muscular inco-ordination is probably the most feasible. It can occur at any age and usually commences with some swallowing difficulty which increases until the patient is only able to imbibe fluids. Often, patients are not wasted in spite of being unable to swallow solid food. As they are forced to take fluids alone, they swallow quantities of milk and highly nourishing liquids. As a result of the delay in emptying, the oesophagus becomes patulous and is often large enough to contain several pints of fluid. The oesophagus may over-fill and its contents may run into the trachea with a resultant broncho-pneumonia (Figs. 5A, 5B).

The onset of symptoms is so gradual that a few years may elapse before the patient seeks medical aid.




Fig. 5A. An early stage of achalasia of the cardia causing dysphagia, which responded well to oesophageal dilatation.

Fig. 5B. A case of achalasia of the cardia of several years' duration. The widely distended oesophagus is seen as a broad mediastinal shadow.

3. *Peptic Oesophagitis*. This lesion is due to the presence of secretory glands in the lower end of the oesophagus. Peptic ulceration occurs in this abnormally placed tissue in similar fashion to the ulceration seen in the gastric mucosa.

With the passage of time the inflammatory reaction is supplanted by the formation of fibrous tissue which undergoes contraction. The resulting stricture is responsible for the obstructive symptoms.

4. *Cicatricial Stenosis*. This is a condition which frequently complicates a hiatus hernia and short oesophagus. In these cases there is ulceration at the cardia and the fibrous contraction causes not only a stricture effect, but also an apparent shortening of the oesophagus.

5. *Diaphragmatic Hernia*. Herniation of part of the stomach through the diaphragm leads to a distortion of the lower end of the oesophagus and also to its compression by the additional bulk contained in the posterior mediastinum.

Some degree of oesophagitis is present in most cases of old-standing diaphragmatic hernia. Stricture of the oesophagus may be a sequel, with narrowing of the oesophageal lumen as well as shortening of the tube. This shortening has often been considered congenital in origin. Congenital shortness of the oesophagus occurs but rarely, however, and in

cases where a diaphragmatic hernia is present, the true position should be readily ascertained (Fig. 6).

6. *Traction Diverticulum*. A diverticulum may result from the adherence of the oesophagus to some localized site of infection, e.g. an infected tuberculous gland. Peristaltic movements will lead to an increase in the size of a hernia once the muscular wall of the oesophagus has been weakened. The mechanism of obstruction is similar to that seen in the case of a pharyngeal diverticulum. The food passes straight into the diverticulum or blind end, and as this enlarges the part of the oesophagus below the sac comes off at a higher and more oblique angle. The outlet from the diverticulum may become obstructed, or the actual weight and pressure of the diverticulum on the lower oesophagus may cause dysphagia.

E. OTHER CAUSES

Several unusual causes of dysphagia should be mentioned for the sake of completeness:

- (a) Simple tumours.
- (b) Sarcomata.
- (c) Syphilitic stricture.
- (d) Scleroderma.

ASSOCIATED SYMPTOMS

Obstruction. The course of obstruction is usually slow but progressive. Initially the patient may complain of a sensation of something in the chest, which he usually describes as a 'lump'. After a time a difficulty in the swallowing of solid and coarse food is noticed; this progresses until food will only pass down the oesophagus if helped along with a gulp of water.

Eventually only fluids can pass and, at this stage, vomiting or regurgitation may become troublesome. If, perchance, a small portion of food should block an already narrowed oesophagus, complete obstruction will result.

Vomiting. Bleeding associated with a regurgitation of food is suggestive of a carcinoma. A massive growth may not only obstruct the lumen, but may bleed into it. The regurgitation of food and blood may come right up into the mouth.

Anorexia and Loss of Weight. Loss of weight naturally occurs as the general nutrition of the patient regresses. This may be due to a loss of appetite as well as to the actual lack of food passing into the stomach.

Pain and Dyspepsia. Pain is not a common symptom except when a foreign body has been swallowed. Patients frequently complain that they have the sensation of a lump; this is



Fig. 6. Diaphragmatic hernia showing part of the stomach in the thorax. The oesophagus is normal in length, and the hernia appeared only in the Trendelenberg position.

usually referred to the centre of the chest. A drink of water may be required to facilitate the passage of foods which feel 'stuck in the chest'. Discomfort following food or discomfort experienced between meals is frequently ascribed to the burning of indigestion.

GENERAL EXAMINATION

The general examination must be a complete one and should include particularly careful inspection of the mouth, tongue and throat, and an examination of the central nervous system.

It is not enough to listen to the patient's story of a swallowing difficulty. He should be asked to demonstrate it, as during the process of swallowing, auscultation of the chest may indicate the site at which the passage of the food is obstructed.

Vomited material should be inspected. There may be no complaints of melaena, but the patient's stools should always be examined as a slight bleeding may have been overlooked.

The patient should be weighed. Strangely enough, he is often totally unaware of a loss of weight. As mentioned before, patients suffering from dysphagia may actually be obese, because they drink rich milk foods when they are unable to swallow more solid material. They thus have a much greater total food intake than the average person.

The skin should be examined for changes due to scleroderma or to hyperthyroidism.

OPSOMMING

Die etiologie van disfasie word bespreek, met spesiale verwysing na karsinoom van die slukderm.

(To be continued)

SWELLINGS OF THE NECK

MIDLINE SWELLINGS

JAN H. LOUW, CH.M.*
and

C. F. M. SAINT, C.B.E., M.D., M.S., F.R.C.S., Hon. F.R.A.C.S., Hon. F.R.S.M.†

Department of Surgery, Medical School, Mowbray, C.P.

(Continued from p. 288)

2. INFLAMMATORY SWELLINGS

Although almost any inflammatory swelling may occupy the midline of the neck, the only infective lesion which is *characteristically* midline is the gumma of tertiary syphilis. Midline gummata of the neck are usually found below the level of the thyroid cartilage in the suprasternal region. They may be limited to the soft tissues (Fig. 11), but more commonly start as a sub-periosteal focus which subsequently extends to the overlying skin (Fig. 12).

A gumma is a chronic granulomatous swelling consisting of a central mass of necrotic tissue surrounded by a zone of cellular infiltration and fibrosis. It commences gradually and indefinitely with the development of a comparatively painless, rounded or ovoid indurated swelling, the edges of which fade into the surrounding tissues. If untreated, the mass enlarges slowly and after weeks or even months softens at the centre. The swelling



Fig. 11 (Left). Midline gumma of the skin in the suprasternal region.

Fig. 12 (Right). Subperiosteal gumma of the sternum which has broken down to form a typical gummatous ulcer.

is at first firm and fibrous; later it becomes 'rubbery' and finally soft and fluctuant. When the inflammatory process approaches the surface, the overlying skin becomes hyperaemic and assumes a purplish colour.

Individual gummata seldom exceed the dimen-

* Professor of Surgery, University of Cape Town.
† Emeritus Professor of Surgery, University of Cape Town.

sions of a small orange, at which stage the swelling breaks down on the surface discharging thin, yellowish pus. A typical gummatous ulcer with punched-out edges and covered by a wash-leather slough is left (Fig. 12).

The diagnosis of a gumma should not present much difficulty. Syphilis must be suspected in any comparatively painless swelling of obscure origin. If such a swelling has the features of a mild, chronic inflammatory process and there are other manifestations of syphilis, the diagnosis is straightforward. Serological tests will provide confirmatory evidence and, under appropriate treatment, the lesion usually resolves with surprising rapidity.

3. MEDIAN LINE SWELLINGS OF THE THYROID

Thyroid swellings usually extend well beyond the midline of the neck and will be dealt with under the heading of *Swellings of the Anterior Triangle*. In certain circumstances, however, localized nodules in the thyroid present as midline swellings to be differentiated from other lumps in this region. The following examples illustrate this.

(a) *Adenoma of the Isthmus* (Fig. 13). Adenomata of the thyroid arising from the isthmus may either bulge forwards between the lower bellies of the sternomastoids, or descend behind the sternum. The latter tend to produce pressure on the trachea and the patients often complain that bending of their neck forwards, or elevation of their arms,

causes choking, especially at night.

These adenomata, like adenomata arising in other parts of the gland, are usually single, and the rest of the gland is perfectly normal. Symptoms of thyrotoxicosis are rare, but evidence of malignancy may be present. Whenever such a discrete nodule is found in the thyroid, its immediate removal must be advised because of the risk of malignancy. (This will be discussed in a later section).

(b) *Adenoma of the Pyramidal Lobe* (Fig. 14). It is not generally appreciated that a pyramidal lobe is present in fully 70% of people, and occasionally an adenoma may develop in it. Such an adenoma may simulate a thyroglossal cyst very closely, especially if it has undergone cystic degeneration. As a rule, however, a tongue of thyroid tissue can be felt either above or below the adenoma. Immediate removal is the only rational treatment.

(c) *Hyperplasia of the Pyramidal Lobe* (Fig. 15). Unless the pyramidal lobe is carefully identified and removed during the course of a thyroidectomy, a portion of this little tongue of thyroid tissue may inadvertently be left *in situ*. We have encountered a number of these cases where the remnant subsequently enlarged with the production of a swelling resembling a thyroglossal cyst. The presence of a thyroidectomy scar and, in one patient, the recurrence of thyrotoxicosis, rendered the diagnosis of our cases straightforward.



Fig. 13 (Left). Solitary adenoma of the isthmus of the thyroid.

Fig. 14 (Middle). Solitary adenoma of the pyramidal lobe of the thyroid.

Fig. 15 (Right). Hyperplasia of the pyramidal lobe. The patient had a previous thyroidectomy (note scar just above clavicles) and this swelling arose in a remnant of pyramidal lobe not removed.



Fig. 16 (Left). Tuberculous submental glands presenting as a paramedian swelling.

Fig. 17 (Middle). Suppurative submental lymphadenitis producing a midline swelling simulating an infected dermoid cyst.

Fig. 18 (Right). Suppurative lymphadenitis simulating an infected thyroglossal cyst.

B. OTHER MIDLINE SWELLINGS

These include paramedian swellings as well as swellings which have extended across the midline or happen to be midline simply by chance. They arise mainly in the skin, subcutaneous tissues, lymph nodes and thyroid gland and may be due to congenital malformations, trauma, inflammation or neoplastic formation. All these swellings are more fre-

quently encountered in the lateral triangles of the neck and will be considered in greater detail when lesions of those regions are discussed. Reference is being made to them at this stage because in this situation they may be confused with typically midline swellings. Illustrative examples that we have encountered include the following:

1. *Lymph Node Affections.* It has already been pointed out that no single lymph node is situated strictly in the midline of the neck. Many of them, however, are situated close to the midline, e.g. submental, cricothyroid and suprasternal. When these enlarge the swelling may simulate a midline swelling very closely (Fig. 16). Indeed, if glands on either side of the midline enlarge and become adherent, a truly midline swelling is produced.

If suppuration should occur in these enlarged glands, the abscess may spread in such a way that it occupies the midline and simulates an infected dermoid (Fig. 17) or a thyroglossal cyst (Fig. 18).

Furthermore, a cold abscess arising in tuberculous glands may penetrate the deep fascia and point in the midline (Figs. 19 and 20) or it may present in the suprasternal fossa (Fig. 21).

Malignant lymph node enlargements may also present as midline swellings. Examples which we have encountered include secondaries from carcinoma of the lip (this Journal, 1955, p. 102, Fig. 36), secondary melanoma (this Journal, 1955, p. 106), secondaries from carcinoma of the breast simulating a thyroid swelling (Fig. 22) and Hodgkin's lymphadenopathy (Fig. 23).




Fig. 19. "Collar-stud abscess" arising from tuberculous cricothyroid glands and presenting in the midline.



Fig. 20 (Left and middle). Tuberculous "collar-stud abscess" simulating an infected thyroglossal cyst.
Fig. 21 (Right). Cold abscess presenting in the suprasternal fossa.



Fig. 22 (Left). Secondary suprasternal malignant glands from carcinoma of the breast, simulating a thyroid swelling.

Fig. 23 (Right). Hodgkin's disease of the cervical lymph nodes.

The clinical features of all these glandular enlargements have already been described. Their diagnosis does not usually offer much difficulty.

OPSOMMING

Ontstekingswelsels en middel-lyn-swelsels van die skildklier word geïllustreer en beskryf, sowel as ander swelsels wat uit 'n kliniese oogpunt na middel-lyn-swelsels lyk, nl. limfklieraandoenings.

(To be continued)

THE HYPERVENTILATION OR OVERBREATHING SYNDROME

FRANCES AMES, M.D.

Department of Neuropsychiatry, Groote Schuur Hospital, Observatory, C.P.

Most people breathe too fast or too deeply when they are afraid or angry or emotionally disturbed in some other way. The reason for this increase in rate or depth of breathing is not known. It may be part of a primitive preparation for fight or flight. Cannon suggested that it served the useful purpose of getting rid of carbon dioxide from the body in anticipation of the augmented discharge of carbon dioxide from the muscles as soon as the great muscular exertion required for fight or flight begins. Primitive man had no hesitation about the necessity for muscular effort when aroused. Modern man is, to a large extent, inhibited by cultural and social tradition from expressing his anger or fear so directly. Consequently, although physiologically prepared for such activity, he has no outlet for it in muscular exertion. The overbreathing may then lead to a distressing group of symptoms and signs which have been labelled the 'hyperventilation syndrome'.

Those who suffer from this syndrome seldom complain of overbreathing. Their subjective respiratory difficulty has been called 'sighing dyspnoea'. They usually describe it as an inability to 'take a deep enough breath' or 'get enough air into the lungs'. It can be an extraordinarily distressing symptom. One patient drove through town holding the door of his car open because of his urgent desire to get more air into his lungs. Another patient who usually got his attacks at night (nocturnal occurrence of attacks that wake the patient from sleep are not uncommon) would jump out of bed twisting and writhing in his efforts to 'get a deep enough breath'.

A striking feature of this respiratory difficulty is that it is never related directly to exertion. Indeed, in most cases it is relieved by motor activity and patients often remark on their inability to remain still during an attack. Relatives are often aware of the overbreathing but may omit to mention it because their attention is directed to other symptoms or signs. One patient had been diagnosed as epileptic although her 'fits' were clearly hyperventilation attacks. No mention was made of any respiratory difficulty until her mother watched her being instructed to overbreathe

and said: 'Oh, but she always breathes like that before she gets a fit'.

Overbreathing causes marked temporary disturbance of neurogenic function, both cerebral and peripheral. More than 60% of people who overbreathe complain of a feeling of light-headedness or giddiness—never a true vertigo. This sensation is aggravated by the upright posture and relieved by recumbency. Most patients feel unsteady on their feet and may appear to be intoxicated. One woman of 45 had noticed that if she got bad news, she tended to have difficulty in 'getting enough air into her lungs'. On one occasion she had an attack of this sort while travelling on a bus. She felt confused and her extremities began to tingle. She felt so unwell that she asked the conductor to stop the bus so that she could dismount. She spoke thickly because her mouth felt stiff and she staggered as she spoke. He thought she was drunk and refused to stop the bus. In desperation she jumped off while it was still moving and he threatened to charge her for drunk and disorderly conduct.

Some patients describe their cerebral condition during hyperventilation as being one of a most disturbing 'unreality'. A doctor once described to me a seance which he attended in the company of a young girl who had recently been bereaved. He said: 'In the darkness I could hear her breathing heavily. Suddenly she screamed and as the lights were switched on and a pencil thrust into her hand she scrawled DAD with tetanic fingers.' The girl, because of the strange feeling of unreality induced by overbreathing, was quite sure she had been in touch with the spirit of her dead father.

The term 'black-out' was frequently used by patients. Of a series of 40 cases only 5 had a disturbance of consciousness gross enough to merit the term. A typical case is the schoolteacher who was sitting reading a book when she suddenly felt faint and began to sweat. She then became unconscious within a few minutes of the onset of her symptoms and said that she was completely unaware of her surroundings for several minutes. When she came to she noticed that her thumbs were

turned into her palms and her fingers were blue and tingling. She was admitted to hospital as a 'coma of unknown origin'.

Drowsiness is not uncommon. Some persons voluntarily overbreathe in an effort to fall asleep.

Dysfunction of the peripheral nerves leads ultimately to tetany. Sensory involvement is earlier and more common than the well-known motor manifestation of tetany. Paraesthesiae always start distally either in limbs or round the centre of the face. In addition to tetanic spasm of the extremities, difficulty is often experienced in articulation because of 'stiffness' of the muscles round the mouth. Neck rigidity may result from spasm of neck muscles. Three patients were sent in to hospital as cases of meningitis because of this sign.

Marked muscle tenderness may be present in muscles previously the site of tetanic spasm. One nurse was labelled 'poliomyelitis' because of tenderness of her calf muscles after a bout of overbreathing.

Apart from the nervous system, other widespread disturbances may occur. The cardiovascular system is disturbed. Tachycardia is an inevitable accompaniment of overbreathing and praecordial pain of the left mammary type is not uncommon. Some authors have described electrocardiographic changes (mainly inversion of T waves) and this may lead the unwary into making a diagnosis of coronary thrombosis in an already anxious patient.

The most common gastro-intestinal disturbance is a feeling of fullness or distension due to concomitant air swallowing. Cardiospasm sometimes results and may cause severe sub-sternal pain.

All patients who overbreathe complain of fatigue 'as though I have just run a mile'. Finally, a striking peripheral vasoconstriction is invariably present.

The pathogenesis of all these symptoms and signs is of interest. At least 3 separate physiological mechanisms are invoked as a direct result of overbreathing. These are:

1. Increased alkalinity of the blood.
2. Reflex peripheral vasoconstriction.
3. Circulatory effects resulting from the muscular exercise of overbreathing.

1. The increased alkalinity of the blood which results from the loss of CO_2 during hyperventilation is of primary importance. Patients who overbreathe a CO_2 -rich mixture do not develop the characteristic symptoms of neurogenic dysfunction. Conversely, many of the symptoms may be relieved by re-breathing CO_2 . It is the alkalosis that:

- (a) Increases neuro-muscular irritability to the

point where tetany may develop.

(b) Causes cerebral blood vessels to constrict. This conserves their low CO_2 content. Cerebral blood flow is usually diminished by about 30% during hyperventilation.

(c) Renders haemoglobin less able to part with its oxygen, thereby interfering further with the nutrition of the tissues (and especially of nerve tissues).

2. Reflex peripheral vasoconstriction is an integral part of the hyperventilation syndrome. The motor route for this reflex is along the sympathetic nerve fibres. The afferent path is not precisely known, but seems to be connected in some way with expansion of the chest or lungs. The peripheral vasoconstriction appears to be of some additional importance in the genesis of the peripheral nerve dysfunction by lowering the threshold of nerve irritability and thus rendering it more susceptible to the effects of the alkalaemia.

3. Hyperventilation is a vigorous form of muscular exercise so that an increase of heart rate occurs regularly.

4. A fourth factor which must receive mention because of the role of emotional stress in the etiology of the hyperventilation syndrome is adrenaline release. It must be considered as a possible background factor because of its action as a respiratory stimulant, its effect in causing skin vasoconstriction and its influence in disturbing ionic equilibrium. Although adrenaline release does not appear to alter the symptoms and signs of the hyperventilation syndrome in a qualitative sense, it plays a part in enhancing or maintaining the physiological effects of the other 3 factors.

TREATMENT

Much of the treatment is built around the hyperventilation test, which consists of instructing the patient to breathe deeply at a rate of between 30-40 breaths a minute. Cerebral symptoms are more easily elicited with the patient upright and peripheral symptoms with the patient recumbent. Within a few minutes the patient's symptoms are reproduced and the recognition of the similarity of the symptoms is immediately and dramatically demonstrated. An explanation of the mechanisms of the hyperventilation syndrome is then given and the fact that emotional stress causes overbreathing is emphasized. The relief gained by re-breathing CO_2 from any convenient receptacle, e.g. a paper bag or hat, is demonstrated.

Finally the patient is encouraged to discuss his emotional problems. Mild sedation may be necessary and the art of progressive relaxation can be taught.

OPSOMMING

Hiperventilasie waarvan die pasiënt onbewus is, kan 'n verskeidenheid van simptome tot gevolg hê wat tans as 'n entiteit in die 'Hiperventilasie-sindroom' herken word.

Daar kan 'n opvallende tydelike versteuring van neurogeniese funksie, sowel serebraal as randstandig, wees wat aanleiding tot bewusteloosheid en tetanie kan gee.

Die kardiiovaskulêre en die spysverteringskanaalstelsel kan ook versteur word.

Die meganismes wat ten grondslag lê van die produksie van die simptome van hierdie sindroom word bespreek, en die behandeling van die toestand word kortliks omskryf. Aandag word gevestig op die noodsaaklikheid dat die pasiënt sy emosionele probleme moet bespreek.

PREPARATIONS AND APPLIANCES

SIGMAGEN TABLETS

Description: Each *Sigmagen* Tablet contains 0.75 mg. prednisone, 325 mg. acetylsalicylic acid, 20 mg. ascorbic acid and 75 mg. aluminium hydroxide.

Advantages: *Sigmagen* combines prednisone (*Meticorten*), aspirin and ascorbic acid to provide potentiated anti-rheumatic action with analgesic effect and vitamin C stress support. Containing *Meticorten*, which has 3.5 times the therapeutic effectiveness of cortisone or hydrocortisone orally with substantially greater freedom from undesirable activity, *Sigmagen* affords safe reliable corticosteroid therapy, especially designed for a wide range of non-specific rheumatic diseases. The simultaneous



complementary action of the combined anti-rheumatic agents in *Sigmagen* permits maintenance of superior relief at minimal dosages unlikely to elicit undesirable effects.

Sigmagen is of particular value in patients no longer responding to salicylates alone.

Indications: *Sigmagen* is indicated in the treatment of mild cases of rheumatoid arthritis or spondylitis, subacute or interval gout, bursitis, myositis, synovitis, fibrositis and neuritis.

Packing: *Sigmagen* Tablets, bottles of 30, 100 and 500.

Manufacturers: Scherag (Proprietary) Ltd., P.O. Box 7539, Johannesburg.

DELTON COMPOUND

SALICYLAMIDE TABLETS

A compound tablet of the antipyretic-analgesic type, based on salicylamide in balanced combination with phenacetin, codeine phosphate and caffeine.

Each Delton tablet contains:

Salicylamide	54.16%
Phenacetin	8.33%
Codeine Phosphate	0.91%
Caffeine	8.33%

Clinical Experience: Salicylamide is a modified form of aspirin. It is a central analgesic of the antipyretic-analgesic type. It is quick in action and less likely than aspirin or salicylate to give rise to gastric irritation. It does not accumulate. In the amounts equal to the usual dosage of aspirin taken to alleviate headache and various aches and pains (5-10 grains) toxicity is extremely improbable. As to the rare case of idiosyncrasy:

(a) There is no evidence that salicylamide is more likely to give rise to excessive reactions than any other substance of this or any other type.

(b) Weight for weight, it is less toxic than aspirin.

There is so far no evidence of allergic manifestations. These are by no means unknown with aspirin, but the lower plasma levels of the amide should tend to make such an unfavourable response less likely with it.

Competent reports from expert physicians, dealing with several hundreds of patients, indicate that in massive dosage it has a better therapeutic index than salicylate or aspirin. Idiosyncrasy should always be probed by administration of a prior test dose before any form of intensive therapy and is the physician's responsibility.

The substitution of salicylamide for aspirin in comparable doses would be an advantage. This substitution in various compound tablets is, therefore, to be recommended since salicylamide has been shown to be compatible and synergistic with phenacetin, codeine phosphate, etc. in addition to its advantages over aspirin.

Indications: Delton can be administered in all cases in which salicylate therapy is indicated. These include acute rheumatic fever, rheumatoid arthritis, fibrositis and polyarthritis. Delton also relieves sciatica, neuritis, myalgia, lumbago, pleurisy, neuralgia and sinusitis.

Dosage: For headaches, neuralgia, toothache, periodic pain—one tablet. Repeat 2-3 times a day, if needed.

For fibrositis, sciatica, lumbago, neuritis or muscular rheumatism—two tablets. Repeat 1-2 times a day, if needed.

For influenza, feverish colds—two tablets, taken with a hot drink, before going to bed.

Administration: Delton tablets can be taken whole, or dispersed in water or milk. They should not be given to children under five. For older children, half to one tablet, according to age.

Packing: Delton is presented in 11.8 gr. tablets and packed in glass phials of 10 tablets, protected by a carton.

Distributors: Protea Pharmaceuticals Limited, P.O. Box 7793, Johannesburg.

PREPARATE EN TOESTELLE

SIGMAGEN-TABLETTE

Beskrywing: Iedere Sigmagen-tablet bevat 0.75 mg. prednison, 325 mg. aetielsalisiezuur, 20 mg. askorbiensuur en 75 mg. aluminiumhidroksied.

Voordede: Prednison (Meticorten), aspirien en askorbiensuur is verenig in Sigmagen om 'n versterkte rumatiekbetrydende uitwerking gepaard met 'n pynverdwende effek en vitamien-C-spanningsteun te verskaf. Sigmagen bevat Meticorten wat terapeuties 3-5 keer doeltreffender is as mondelinge kortison of hidrokortison, en 'n aansienlik groter vryheid van onwenslike aktiwiteit bied. Sigmagen is dus 'n veilige en betroubare kortikosteroid-middel wat spesiaal ontwerp is vir 'n groot verskeidenheid van nie-spesifieke rumatiekkwale. Die gelyktydige aanvullende uitwerking van die gesamentlike rumatiekbetrydende middels in Sig-



magen maak dit moontlik om voortreflike verligting te verskaf met minimale dosisse wat hoogs waarskynlik geen ongunstige effek sal hê nie.

Sigmagen is veral van groot waarde vir pasiënte wat nie langer op salisilate alleen reageer nie.

Indikasies: Sigmagen word aangedui vir die behandeling van ligte gevalle van chroniese gewrigsontsteking of werwelontsteking, subakute of tussenpoosjig, slymbeursontsteking, miositis, gewrigsvliesontsteking, fibrositis en senuwee-ontsteking.

Verpakking: Sigmagen-tablette, bottels van 30, 100 en 500.

Fabrikante: Scherag (Proprietary) Ltd., Posbus 7539, Johannesburg.

DELTON-SAMESTELLING

SALISIELAMIED-TABLETTE

'n Saamgestelde tablet van die koorswerende en pynstillende tipe, gebaseer op salisielamied in 'n gebalanseerde samestelling met fenasetien, kodeienfosfaat en kaffeien.

Iedere Delton-tablet bevat:

Salisielamied	54.16%
Fenasetien	8.33%
Kodeienfosfaat	0.91%
Kaffeien	8.33%

Kliniese Ondervinding: Salisielamied is 'n gewysigde vorm van asperien. Dit is 'n sentrale pynstillende van die koorswerende en pynstillende tipe. Dit werk vinnig, en sal waarskynlik nie so maklik soos asperien of salisilaat aanleiding tot maagprikkeling gee nie. Dit hoop nie op nie. In hoeveelhede gelykstaande aan die dosis asperien wat gewoonlik geneem word om hoofpyn en verskillende ander pyn te verlig (5-10 grein) is toksisiteit hoogs onwaarskynlik. Wat betref die uitsonderlike gevalle van idiosinkrasie:

(a) Is daar geen bewyse dat salisielamied waarskynliker aanleiding sal gee tot buitensporige reaksies as enige ander middel van hierdie of enige ander tipe nie.

(b) Is dit, gewig vir gewig, minder toksies as asperien.

Daar is geen bewyse van allergiese manifestasies nie. Sulke manifestasies is geensins onbekend met asperien nie, maar die laer plasma-peil van die amied behoort so 'n ongunstige reaksie minder waarskynlik te maak.

Gesaghebbende verslae van deskundige geneesheer wat etlike honderde pasiënte daarmee behandel het, dui aan dat, in massiewe dosisse, dit 'n beter terapeutiese indeks as salisilaat of asperien het. Idiosinkrasie moet altyd vasgestel word deur die toediening van 'n voorafgaande toetsdosis voordat enige vorm van intensiewe terapie toegepas word, en dit is 'n verantwoordelikheid wat by die geneesheer berus.

Die vervanging van asperien met salisielamied in vergelykbare dosisse sal derhalwe voordelig wees. Sodanige vervanging in verskillende saamgestelde tablette word derhalwe aanbeveel, aangesien salisielamied die bewys gelewer het dat dit vermengbaar en sinergies met fenasetien, kodeienfosfaat, ens., is, heeltemal afgesien van sy ander voordede in vergelyking met asperien.

Indikasies: Delton kan toegedien word in alle gevalle waar salisilaatterapie aangedui word. Dit sluit in rumatiekkoors, chroniese gewrigsontsteking, fibrositis en poliartiritis. Delton verlig ook heupjig, senuwee-ontsteking, mialgie, lendejig, borsvliesontsteking, sinkings en sinusitis.

Dosis: Hoofpyn, sinkings, tandpyn, periodieke pyn—een tablet. Herhaal 2-3 keer per dag, indien nodig.

Fibrositis, heupjig, lendejig, senuwee-ontsteking of spierrumatiek—twee tablette. Herhaal 1-2 keer per dag, indien nodig.

Influenza, koorsagtige verkoues—twee tablette met 'n warm drankie voor slapenstyd.

Toediening: Delton-tablette kan heel geneem of in water of melk opgelos word. *Hulle moet nie aan kinders onder vyf jaar gegee word nie.* Vir ouer kinders: 'n Halwe tot een tablet, volgens die

ouderdom van die kind.

Verpakking: Delton word aangebied in tablette van 11.8 grein in glaslessies bevatende 10 tablette en beskerm in 'n kartondosie.

Verspreiders: Protea Pharmaceuticals Limited, Posbus 7793, Johannesburg.

NOTES AND NEWS • BERIGTE

DISPENSING BY DOCTORS

The Minister of Health decided not to proceed with the highly contentious Medical, Dental and Pharmacy Amendment Bill during the last session of Parliament. This arrests temporarily the public agitation by pharmacists to prevent doctors from dispensing.

The serious situation created by this attack on the medical profession's rights must be watched with great care during the Parliamentary recess.

* * * *

Dr. Louis F. Freed, M.A., M.D., D.Phil., author of *The Problem of European Prostitution in Johannesburg*, has been invited to attend the forthcoming Congress of the International Abolitionist Federation, to be held at Frankfurt-am-Main from 17-19 October 1956. Dr. Freed is Lecturer on Social Medicine at the University of the Witwatersrand.

* * * *

Mr. Bertram L. Shaff, M.B., F.R.C.S. (Edin.), has joined Mr. J. Wolfowitz, M.B., F.R.C.S. (Edin.) in partnership in practice as a surgeon at 308 Medical Centre, Jeppe Street, Johannesburg. (Telephones: Rooms: 23-5311/2; Residence: 41-2026).

* * * *

CORONARY THROMBOSIS AND AIR TRAVEL

The great majority of patients who have recovered from coronary thrombosis are fit to make a short flight of under 5 hours' duration. But those with even a minor degree of coronary disease should not undertake long journeys by air unless they have been free from symptoms for at least 3, and preferably 6, months. Their condition should be revealed to the medical department of the airline for a decision regarding advisability to travel and to enable appropriate instructions to be issued to all concerned for their care and welfare along the route. (Whittingham, Sir H. (1956): *The Practitioner*, 176, 179.)

REVIEWS OF BOOKS

JOHNSTONE'S MIDWIFERY

A Text-Book of Midwifery for Students and Practitioners. By R. W. Johnstone, C.B.E., M.A., M.D., Hon. LL.D., F.R.C.S.E., F.R.C.O.G., F.R.S.E. (Pp. 572 + Index. With 295 illustrations. 36s.) 1955. 16th ed. London: A. & C. Black Limited.

The sixteenth edition of this well-known and standard textbook of midwifery has been revised in collaboration with Prof. R. J. Kellar, the present incumbent of the Chair of Obstetrics and Gynaecology which Professor Johnstone filled with such distinction in Edinburgh for so many years.

The whole text of the previous edition has been

revised carefully by both authors, considerable alterations having been introduced in the sections dealing with the functions of the placenta, pre-eclampsia, uterine inertia and rupture, and inversion of the uterus.

The first section of the work is devoted to the relevant anatomy and physiology, presented from the standpoint of the clinical implications. The 'applied' approach in the endocrine description, e.g. is well illustrated by the explanation of premenstrual tension and the treatment of the condition based on experimental work, viz. the limitation of fluid intake during the second half of the menstrual cycle in those cases where the physio-



Dr. R. Singer

Dr. Ronald Singer, of the Department of Anatomy, University of Cape Town, has been elected a Fellow of the American Anthropological Association. He

has also been invited to read a paper at the Neanderthal Centenary in Dusseldorf in August 1956.

The Wenner-Gren Foundation for Anthropological Research, New York, recently awarded him a bursary to enable him to carry out specific studies at various research institutions in Europe, Scandinavia and the U.S.A., as well as to facilitate his attendance to read papers at the *First International Congress on Human Genetics* in Copenhagen (1-6 August) and the *Fifth International Congress on Anthropological and Ethnological Sciences* in Philadelphia (1-9 September).

* * * *

FIRST INTERNATIONAL CANCER CYTOLOGY CONGRESS

This important convention, sponsored jointly by the International Union against Cancer, the College of American Pathologists, the American Society for Clinical Pathologists and the Inter-Society Cytology Council, will be held at the Drake Hotel, Chicago, Illinois, U.S.A., from 9-11 October 1956.

An extensive programme, including panel discussions, has been arranged.

Further information can be obtained from Prof. J. Gillman, Department of Physiology, Medical School, Hospital Street, Johannesburg. (Telephone: 44-1492).

* * * *

Mr. Edward Abro, M.B., Ch.B., Ch.M. (Rand.), has commenced practice as a neurologist at 609 Medical Centre, 209 Jeppe Street, Johannesburg. (Telephones: Rooms: 22-1214; Residence: 41-4695).

logical disturbance of the electrolyte pattern creates a condition which borders on the pathological.

Very clear and ample diagrams are a feature of the book. They illustrate admirably the account of normal and abnormal pregnancy and labour, and the concise classified presentation of aetiology, differential diagnosis, etc. throughout the work makes this volume a very popular one amongst undergraduate students preparing for their examinations. The important chapter on *Radiography in Obstetrical Diagnosis* (by Dr. J. Z. Walker) is illustrated with excellently reproduced radiographs and diagrams and, apart from operative obstetrics, there are special chapters on the puerperium, the physiology and care of the new born child and its pathology. The latter chapter includes a simplified account of the genetics of Rh with adequate information (from the clinical standpoint) about the Rh antibodies, for which the Fisher terminology is employed. However, the statement (p. 485) that the Rh genes occur at 3 adjacent loci on the chromosome is strongly contested in certain quarters. Even the Fisher terminology requires a fourth locus to accommodate Ff in addition to CcDdEe.

There is a clear statement of the indications for abortion, the view being expressed that, except when it is done with a definite therapeutic aim, abortion is illegal in all Christian countries. South Africa presumably qualifies for inclusion in this category of countries. At any rate, it has made provision for statutory recognition of the Almighty. Yet it is doubtful whether a therapeutic aim not directed to the prevention of an imminent threat to the life (as opposed to the health) of the mother would be regarded as lawful in the Union, where we have in addition such serious problems as pregnancy resulting from rape, especially when White and Black races are involved.

It is of interest that the authors regard German measles in the early months of pregnancy as a legitimate indication for abortion (p. 540), since in such a case there is no therapeutic aim directed at the mother; on the contrary, the intention is to destroy a probably malformed foetus. It is doubtful whether such an indication is lawful in the Union.

Many generations of students have been raised on this excellent textbook and have found it of great value subsequently in practice. The instruction in the guidance given for the management of obstetrical conditions is so sound that it is likely to find fairly world-wide acceptance. Its suitability as a textbook should, therefore, meet a considerable appeal, despite the local idiosyncrasies of teachers of the subject in various countries.

SCIENTIFIC BASIS OF MEDICINE

Lectures on the Scientific Basis of Medicine. Vol. III, 1953-1954. (Pp. 398. With IX plates. 35s.) London: The Athlone Press.

This is the third volume selected from the annual series of lectures given under the auspices of the British Post-Graduate Medical Federation and, like the previous volumes, must prove a source of pleasure to the progressive post-graduate student who looks upon medicine not only as a technical craft and a means of making a living, but also as an intellectual pursuit and a philosophy.

It is difficult to select individual lectures for comment, as the over-all standard is so excellent, but the first lecture by Professor Sigerist on *Science and History* should be read by all medical teachers and specialists. Professor Sigerist points out that

the 'expert' has also 'an important part to play as an expert citizen', and that if the scientist is to take his rightful place in the community at large, the sciences must be approached not only technically but 'also historically, philosophically and sociologically'.

Professor Garrod contributes a lecture on *Causes of Failure in Antibiotic Therapy*, which presents simply and in common-sense terms the difficulties encountered by practitioners, and how to overcome these difficulties. For those lost in the welter of new drugs used in the treatment of malaria, Dr. C. Chesterman's lecture on *Anti-Malarial Drugs* will throw a beam of light into the darkness.

Other lectures on the *Chemotherapy of Cancer*, *Stress and Thyroid Activity*, *Body Water Control*, *The Supporting System and its Disorders*, show the diversity of the subjects covered. Professor Haldane's lecture on *The Genetics of Some Biochemical Abnormalities*, and Dr. G. Popjak's on *Biological Synthesis*, provide food for thought for those who are interested in these scientific border-lines of medicine. Dr. R. MacKenna's *Scientific Approach to Dermatology* will convince the sceptics who do not at present regard dermatology as any more than a matter of clinical duty.

While there is no doubt that some of the lectures will prove too technical for the ordinary reader, the book will prove extremely valuable to all practitioners who like to read beyond the bounds of their own special interests, and like to undergo an annual course of 'brain dusting'.

ELECTROCARDIOGRAPHY

A Primer of Electrocardiography. By George E. Burch, M.D., F.A.C.P. and Travis Winsor, M.D., F.A.C.P. (Pp. 286. With 281 illustrations. 1955. 3rd ed. \$5.00). Philadelphia: Lea & Febiger.

The third edition of this book maintains the standard set by its predecessors. The text deals comprehensively with the fundamental concepts and the complex problems of modern electrocardiography. Hence the use of the word *Primer* in the title may, in a sense, be misleading and has no doubt been used to indicate the authors' aim to stress the basic rather than the contentious ideas in present-day electrocardiography. Controversial matters have been omitted and the essential principles are dealt with in a detailed and extensive manner. This detail may possibly cause a little bewilderment to the uninitiated, but is nevertheless very useful to the cardiologist who has occasion to refer to or who wishes to increase his knowledge of the basic principles.

The text has been profusely illustrated with many meaningful diagrams which contribute much to the understanding of the subject. A clear diagram is invariably worth pages of text.

The material is presented in sections dealing with the analysis and clinical significance of the electrocardiogram, the precordial leads, the disorders of cardiac rhythm and the applications of electrocardiography in clinical medicine.

Perhaps the explanation and interpretation of the various components of the electrocardiogram could have been carried out to a greater extent in terms of unipolar rather than bipolar electrodes. An innovation is an introductory section on spatial vectorcardiography. An excellent Appendix comments on the electrical axis, the tri-axial reference system and the average norms of the various intervals and component waves of the electrocardiogram. Another

section of the Appendix lists the definite and indefinite electrocardiographic criteria for the diagnosis of myocardial disease.

The text is well presented, clearly printed and bound in a handsome cloth cover. It is well suited to the experienced cardiologist as well as those beginners who have already made some acquaintance with the subject.

VIRAL AND RICKETTSIAL DISEASES OF MAN

Viral and Rickettsial Diseases of the Skin, Eye and Mucous Membranes of Man. By Harvey Blank, M.D., and Geoffrey Rake, M.B., B.S. 1955. Pp. 263 + Index. With 36 full colour illustrations and 63 black and white illustrations. 60s.). London: J. & A. Churchill Ltd.

This attractive book should interest not only specialists in diseases of the skin and the eye, but also every practitioner who is confronted daily with diseases of viral causation. After two introductory chapters, which include a practical résumé of methods used for the diagnosis of viral diseases, there are detailed accounts of herpes simplex, chicken pox and zoster, smallpox, cowpox and vaccinia, several exanthemata (including Fourth and Fifth Diseases) the common cold, herpangina, warts and molluscum contagiosum. These are followed by a concise yet adequate account of relevant viral diseases contracted from animals, including milker's nodules, orf, foot and mouth disease, cat scratch disease and psittacosis, and further chapters on epidemic keratoconjunctivitis, trachoma, lymphogranuloma venereum and the rickettsial diseases.

The inclusion of so much material in a comparatively small volume deserves great praise; the authors have nowhere sacrificed clarity to conciseness and every subject receives enough treatment to interest and instruct the practising medical man. A few points of general interest are picked out of the host of instructive material presented by the authors, in the hope that they will stimulate practitioners to read this book.

Contrary to current belief, there is no evidence that recurrent aphthae are caused by a virus, particularly the virus of herpes simplex. Treatment of trachoma with sulphonamides and antibiotics apparently attacks the causative virus (a Chlamydozoacea) directly, and not only the secondary bacterial invaders. 'Complete surgical excision of warts cannot be recommended because at least 25% of the lesions recur in the scar'. This fact is well known to dermatologists, but it is time that it was also revealed to many surgeons—in this country at any rate. The whole section on the treatment of warts is, in fact, a gem of summarization and should prove of great value to the practitioner.

The few minor criticisms that follow do not imply that this reviewer has anything but the highest praise for the book as a whole. The Koebner phenomenon refers to the reaction produced by non-specific irritation imitating the clinical dermatosis; thus a patient with psoriasis may develop a psoriatic stripe at the site of a scratch. It does not refer to the inoculation of virus and subsequent development of warts in a scratch, as inferred on pp. 10 and 170. The use of xylol or clove oil as an aid to the examination of plantar warts is not mentioned. The prevalent confusion between 'suck' and 'suckle' is perpetuated on p. 201: 'Direct infection can also occur, as when a previously infected ewe is suckled by an infected lamb'. South Africans will be pained or flattered (as the case

may be) to observe that G. H. Findlay has been confused with G. M. Findlay in the author index.

The book is attractively produced, with well chosen and well produced illustrations. The colour plates are outstanding and an object lesson which some publishers might well study.

YELLOW FEVER VACCINATION

Yellow Fever Vaccination. By K. C. Smithburn, C. Durieux, R. Koerber, H. A. Penna, G. W. A. Dick, G. Courtois, C. de Souza Manso, G. Stuart and P. H. Bonnel. (1956. Pp. 231 + Index. With Figs. and Tables. 25s.) Geneva: World Health Organization.

First applied in French West Africa in 1934, and then in Brazil in 1937, systematic vaccination against yellow fever has immunized tens of millions of persons, and human yellow fever has become a rare disease. When vaccination was begun, however, not even its most optimistic advocates could have anticipated all the benefits and the enormous economic and social advantages that would accrue to those territories where the disease was rife.

The World Health Organization considered that it would be appropriate at this stage to publish a monograph to which would contribute eminent specialists whose names are linked with the success of yellow fever control the world over.

In the first section Smithburn describes the basis on which yellow fever immunization now rests.

In French West Africa and French Equatorial Africa more than 56 million vaccinations were performed between 1939 and 1953 with the scratch vaccine prepared by the Institut Pasteur at Dakar; these have resulted in a protection rate such that human yellow fever has virtually disappeared from the territories in which vaccination was performed. Durieux, formerly Director of the Institut Pasteur at Dakar, describes the preparation and control of the vaccine produced by that Institute, the technique of its administration, and the system of mass vaccination applied in French African territories. Together with Koerber he discusses the immunological results obtained.

The most widely used vaccine in the rest of the world is 17D vaccine. The Oswaldo Cruz Institute at Rio de Janeiro is one of the most important establishments preparing it. Penna, who directs the Institute's yellow fever laboratory, gives details of the production and administration of 17D vaccine which, like that of the Institut Pasteur at Dakar, is used in mass vaccination campaigns. de Souza Manso reports on what has been accomplished with these campaigns in Latin America.

Courtois records the results of numerous control experiments which clarify previous knowledge of the time of appearance of immunity to yellow fever after vaccination and of the duration of that immunity.

Like all vaccination, that against yellow fever has sometimes been followed by more or less serious reactions. It is necessary to know about these accidents and only proper to draw attention to them. They in no way detract from the value and efficacy of the method. Stuart gives a detailed and complete account of reactions observed to vaccinations with both the Dakar and the 17D vaccines.

Each method of vaccination has its advantages and its disadvantages. Vaccination with 17D vaccine by the scarification method (described by Dick) seems to combine the advantages of both Dakar and 17D vaccines without the possible disadvantages of

the more usual ways of administering either.

Vaccination against yellow fever is officially sanctioned as a preventive method by international health legislation, which allows travellers holding a valid certificate of vaccination to avoid quarantine isolation when passing from an area infected with yellow fever to an area receptive to it. Bonnel describes these international regulations and the conditions under which a valid international certificate may be issued.

The monograph also contains a bibliography of works on yellow fever immunology and vaccination.

INSECT VECTORS AND AIR TRAFFIC

Control of Insect Vectors in International Air Traffic: A Survey of Existing Legislation. (Pp. 60. 3s. 6d.) World Health Organization, Geneva.

The insect vectors of diseases such as yellow fever,

malaria, filariasis, etc., may be conveyed (sometimes in considerable numbers) by aircraft on international flights.

Although the international sanitary regulations contain certain articles designed to control the spread of communicable diseases by insect vectors in international air traffic, local regulations for preventing the introduction of such vectors into their territories have been adopted by a number of countries.

This WHO study, which is an analytical survey of the regulations for the disinsection of aircraft in 98 States and territories, includes sections dealing with disinsection requirements, when disinsection is carried out, methods of disinsection, formulations of insecticides used, dosage and time of application, measures taken to prevent the spread of agricultural pests, as well as a summary of the recommendations on the disinsection of aircraft made to the Organization by its Expert Committees.

CORRESPONDENCE

REHABILITATION ASSOCIATION FOR INJURED WORKMEN

WORKMEN'S REHABILITATION CENTRE

To the Editor: I have much pleasure in advising you that the Rehabilitation Association for Injured Workmen has established the 'Workmen's Rehabilitation Centre' at 15 Esselen Street (Corner King George Street), Hospital Hill, Johannesburg, which provides full hospital treatment and physical rehabilitative facilities for European workmen undergoing medical treatment for injuries sustained in accidents which are acceptable under the Workmen's Compensation Act. This Centre is entirely independent and is not controlled by any hospital, nursing home, private hospital or nursing institution.

The Rehabilitation Association for Injured Workmen is a non-profit company registered under the Companies Act, and representatives of insurance carriers under the Act, including the Workmen's Compensation Commissioner, serve on its Board of Directors.

The Centre has been designed especially to cater for serious traumatic cases which would normally entail lengthy absences from work and/or result in serious permanent disablement, the object being to reduce as far as possible the period of temporary disablement as well as the resultant degree of permanent disablement. In other words, the aim of the Association is to restore the injured worker as nearly as possible to his pre-accident productive capacity as soon as possible. This will operate to the economic and social advantage not only of the worker himself but also of the employer and the community as a whole.

The injured workman has free choice of doctor. Any doctor who wishes to admit a Workmen's Compensation patient to the Centre should contact the Medical Administrator or his Deputy by telephone (44-9101).

When an injured workman admitted to the Centre has not yet selected his own doctor and is not in a fit state to do so, a practitioner will be nominated by the Medical Association of South Africa (Southern Transvaal Branch) from its own panel of doctors, to take charge of the case.

In addition to up-to-date theatre and other facilities, the Centre also provides facilities for:

(a) Physiotherapy;

- (b) Occupational therapy;
- (c) Gymnastic and remedial exercises;
- (d) In-patient and out-patient treatment;
- (e) Socio-economic guidance.

The Association's Secretariat at the Centre will maintain close contact with insurance carriers under the Workmen's Compensation Act in regard to cases admitted to the Centre, and will ensure, *inter alia*, that:

i. Assistance will be given to doctors in maintaining records;

ii. A secretarial service is made available to those medical men who have patients at the Centre, and who wish to dictate notes.

Regular clinical meetings will be arranged by the Medical Administrator to discuss clinical problems brought to the meeting by doctors treating cases at the Centre. These meetings will be attended voluntarily by practitioners, and all doctors are welcome to attend.

No attempt will be made to impose upon any doctor any direction about how he should deal with his case. He will be left complete clinical discretion in handling the problems of the patient. He is entitled to exercise his discretion and choice in consulting any other colleague, including specialists in physical medicine, radiologists, etc.

A medical representative of the Workmen's Compensation Commissioner's office, Pretoria, will be in attendance at these meetings, to assist medical practitioners in medical matters relating to the condition of the injured workman. By this liaison with medical men, the assessment of the degree of disability will be materially facilitated.

The Medical Administrator of the Centre, Dr. Boris Serebro, in an administrative and non-clinical capacity, will be available to supply information, and answer queries on all the aspects and activities of the Centre.

Doctors' accounts should, as in the past, be rendered to the employer concerned, as the Association is not responsible for the payments of medical fees.

Your kind co-operation and support will be much appreciated.

J. P. GROBBELAAR,
Chairman.

Rehabilitation Association for Injured Workmen,
Corner of Esselen and King George Sts.,
Johannesburg.



PARKE DAVIS

have pleasure in announcing the agreement of the Pharmacy Board to the registration of:—

kondig met genoeë aan dat die Aptekersraad toestemming verleen het tot die registrasie van:—

PARKE, DAVIS LABORATORIES (PTY.) LTD.

**as the name of their South
African company**

It will be the continuing endeavour of the company to be of the utmost service to the medical and pharmaceutical professions of Southern Africa.

**as die naam van hulle Suid-
Afrikaanse maatskappy.**

Die maatskappy sal steeds trag om die mediese en farmaseutiese beroepe van Suidelike Afrika tot die beste van hulle vermoë van diens te wees.

PARKE, DAVIS LABORATORIES (PTY.) LTD.

P.O. Box 9971

Posbus 9971

LEISK HOUSE, BREE & RISSIK STREETS, JOHANNESBURG
LEISK HOUSE, BREE- & RISSIKSTRAAT, JOHANNESBURG

Telephone 22-2812

Telefoon 22-2812 Telegramme:

Telegrams: "Cascara", Johannesburg.

„Cascara", Johannesburg.

AND AT PORT ELIZABETH

EN TE PORT ELIZABETH.

Distributors in: Union of South Africa — Lennon Ltd., all branches. Rhodesia & Nyasaland — Ian Wilson (Pvt.) Ltd. Bulawayo. And in Mocambique, Angola, Belgian Congo, etc.

Verspreiders in: Die Unie van Suid-Afrika — Lennon Bpk., alle takke. Rhodesië & Njassaland — Ian Wilson (Pvt.) Ltd., Bulawayo. En in Mosambiek, Angola, Belgiese Kongo, ens.

*The universally prescribed
Vaginal Jelly and Cream
for dependable contraception..*



...with the Ortho Diaphragm

Ortho-Gynol

Ricinoleic acid and
p-Diisobutylphenoxypolyethoxyethanol
in a vegetable gum base
buffered at pH.4.5

Ortho-Creme

Ricinoleic acid and
sodium lauryl sulphate in a cream base
buffered at pH.5.8

Ortho Diaphragm

Coil-spring or flat-spring type in
sizes 55 mm. to 95 mm.



Where the diaphragm is impractical or unacceptable

Preceptin

Vaginel Gel* provides the "barrier" effect in its unique synthetic base. Simply applied with the Ortho Applicator

* Ricinoleic acid and p-Diisobutylphenoxypolyethoxyethanol
in a synthetic base buffered at pH.4.5

LITERATURE ON REQUEST


Ortho Pharmaceutical Limited

HIGH WYCOMBE • BUCKINGHAMSHIRE • ENGLAND



Sole distributors:

ETHICAL PRODUCTS (PTY.) LTD.
Ethical Division of Johnson & Johnson (Pty.) Ltd.
P.O. Box 727 East London



NEW

urine sugar test of unmatched simplicity

'Tes-Tape'

(URINE SUGAR TEST TAPE, LILLY)

'Tes-Tape' completely eliminates the need for test tubes, heat, reagents, or any other paraphernalia in urine sugar determinations. Simply moisten a strip of 'Tes-Tape' with the specimen. After just sixty seconds, compare it with the color chart on the 'Tes-Tape' dispenser. Then read off the percentage of sugar. The selective action of 'Tes-Tape' prevents false positive reactions, assures complete accuracy.

The convenient size of the 'Tes-Tape' dispenser permits you to carry it on house calls for on-the-spot determinations. Your patients also will welcome the convenience, simplicity, and accuracy of 'Tes-Tape.'

Lilly

QUALITY/RESEARCH/INTEGRITY



Convenil REGD.
HOMMEL

**New
Antispasmodic-
Sedative**

'Convenil' is an entirely new synthetic by Hommel. In autonomic nervous system disorders it provides highly effective antispasmodic action with excellent tolerability and freedom from side-effects.

The use of 'Convenil' obviates the hypnotic effects of the common antispasmodics; it is a single-drug prescription; it does not contain any barbiturate.

INDICATIONS

Spastic constipation, menstrual and menopausal disturbance, autonomic dysfunction generally, nervous tension and irritability, travel sensitivity.

FORMULA

Each Dragee:—
Phenylglylactic acid-β-
diethylaminoethyl ester ci:rate — — — 150 mg.

DOSAGE

Adults. 1 Dragee 2 to 4 times daily;
in special cases up to 6 daily.

Children: 1 Dragee 1 to 3 times daily.

PACKS

20's and 250's (dispensing)

TRISAN REGD.
HOMMEL

**Sedative-Expectorant
in
Bronchial Asthma**

Symptomatic treatment of bronchial asthma must include attention to expectoration — mucus plugging is now recognized as a cause of unexpected death—and mental sedation.

In 'Trisan' the two desiderata are effected by classically used drugs: Potassium iodide and Barbitone sodium, the latter reinforced by Chloral hydrate.

Composition

Potassium iodide B.P. 6.03%; Chloral hydrate B.P. 7.11%;
Barbitone sodium B.P. 0.24%;
Alcohol 4.00%; Excipient ad 100%

ACTION

'Trisan' is formulated to reduce spasm frequency, to ease expectoration and to reduce mental upset. *Contra-indicated* in hyperthyroidism.

DOSAGE

4 fl. drachms in water during attacks;
for routine symptomatic management between attacks,
1 to 2 fl. drachms nightly for 2 to 3 weeks.

PACKS

Standard Bottle of 4 fl. oz.
Dispensing Bottle of 16 fl. oz.

PROFESSIONAL SAMPLES & LITERATURE ON REQUEST

HOMMEL PHARMACEUTICALS
121 Norwood Road,
London S.E.24

HOMMEL' ethical range includes —

CONVENIL • DORMUPAX • HAEMATOGEN • HICLOSEEN • HYPERYSIN • NYXOLAN • TRISAN

Our Sole Agents for SOUTH AFRICA: Messrs. LENNON LIMITED, P.O. Box 39, CAPE TOWN. P.O. Box 24, PORT ELIZABETH. P.O. Box 266, DURBAN (NATAL). P.O. Box 928, JOHANNESBURG (TRANSVAAL). P.O. Box 76, EAST LONDON

Sole Agents for SOUTHERN RHODESIA: IAN WILSON (PVT) LTD., P.O. Box 1102, BULAWAYO.



*Concerning
a further advance
in the treatment of
bucco-pharyngeal
INFECTIONS*

Dequadin Lozenges containing a new bactericidal and fungicidal substance are rapidly effective against all the common oral pathogens including those resistant to penicillin. They do not contain any antibiotic or anæsthetic substance.

Dequadin Lozenges are indicated in the treatment of Vincent's angina, tonsillitis, sore throat, stomatitis, pyorrhœa, pharyngitis, aphthous ulcers, thrush and glossitis.

They can be used prophylactically in patients exposed to infection especially after tonsillectomy and dental extractions.

By suppressing monilial growths, Dequadin Lozenges prevent the appearance of black tongue and oral thrush. They are also rapidly effective in the treatment of oral thrush including that due to prolonged antibiotic therapy.

DEQUADIN

LOZENGES

In tubes of 20 lozenges each containing
0.25 mg. decamethylene-bis (4-aminoquinaldinium chloride).

S. Africa Pat. No. 22044

CS6/14—13/D

ALLEN & HANBURY'S (AFRICA) LTD
INCORPORATED IN ENGLAND
CONGELLA ROAD • DURBAN



The sedation is in the sugar coating

These tablets, consisting of our well-known Petervite B Formula, plus Phenobarbitone, provide the benefits of the Vitamin B Complex supplementation as well as Phenobarbitone, which exerts the sedation required for patients complaining of nervousness, fatigue and anorexia. In PHEVITAN, we have departed from the normal method of manufacture by the inclusion of the Phenobarbitone in the sugar coating. By this method of administering the sedation, the patient will receive immediate benefit and a feeling of euphoria is the result. This is so vital in nervous patients. The formula: Each tablet contains: Vitamin B₁, 2 Mgm.; Vitamin B₂, 1.5 Mgm.; Vitamin B₆, 0.25 Mgm.; Nicotinamide, 20 Mgm.; Calc. Pantothenate, 2.5 Mgm.; and Phenobarbitone, 1/4 Gr. The Dosage: One tablet before meals and at bedtime. If desired, the bedtime dose may be increased to three tablets for several evenings until the daytime dosage has taken effect. PHEVITAN TABLETS are supplied in: 40's, 100's, and 500's. Price to the patient: 40's—4/6, 100's—9/6, 500's—42/6.

Manufactured in South Africa by



Established 1842

P.O. Box 38 P.O. Box 1200 113, Umbilo Rd. P.O. Box 1005 P.O. Box 2238 P.O. Box 5785
CAPE TOWN BULAWAYO DURBAN BLOEMFONTEIN O.F.S. SALISBURY JOHANNESBURG



DRAEGER INCUBATOR

- ★ INFECTION FREE ACCOMMODATION
- ★ CONSTANT ATMOSPHERE
- ★ CONSTANT TEMPERATURE
- ★ CONSTANT HUMIDITY

With Bacteria-filters, Oxygen Measuring Apparatus, Scale, Acoustic and Light Warning Devices, etc.

SAFETY & MEDICAL EQUIPMENT CO. (PTY.) LTD.

Head Offices: 4th Floor Cambridge, Corner Kerk and Sauer Sts., JOHANNESBURG.

Telephone: 33-9625

P.O. Box 11344



Branch Offices: **CAPE TOWN:** 402-403 A.A. Buildings, Queen Victoria Street, P.O. Box 3599, Cape Town. Telephone 2-5818

DURBAN: 53 Medical Centre, Field Street. Telephones: 6-0894, 6-3036, 2-6318.

Suppliers of:

Oxygen Tents for Adults and Children, Clinical and Transport Incubators, Iron Lungs, Artificial Respiration and Automatic Resuscitation Apparatus such as, Poliomats, Pulmomats, Pulmotors, St. John Ambulance Oxygen Units and Reanimators, Oxygen and Aerosol Therapy Equipment, Machines for Analgesia and Anaesthesia in Dentistry, Midwifery, Gynaecology and Surgery, Hibernation Units, Operating Tables and Lamps, Sterilising Equipment and Plants, Medical Suction Pumps, Suction Massage Units, Audiometers, Oxytest and B.M.R. Apparatus, Pulsimeters, Ergometers, Disinfection Equipment, Rubber and Instrument Specialities.

D7



For Infant Feeding...

Pelargon

FULL CREAM ACIDIFIED MILK



... is an acidified, full cream, powdered milk with added carbohydrates for the feeding either partially or completely of healthy babies from birth onwards. It is also recommended in cases of vomiting and Marasmus, Skin Disorders, Dyspepsia, Diarrhoea and is specially suitable for the feeding of the premature infant.

For further information and literature write to Nestlé's Medical Service, P.O. Box 1568, Durban.

PERCENTAGE COMPOSITION (Dry Powder)

FAT	PROTEIN	LACTOSE	MINERAL SALTS	SUCROSE (added)	MALTO DEXTRIN (added)	STARCH (pre-cooked)	LACTIC ACID	MOISTURE
17	16.5	23.5	4.3	12.5	12.5	8.5	2.3	3

P-256

A NESTLÉ'S PRODUCT

Rapid relief of ASTHMA

with
**BROVON
 INHALANT**

.....

The synergistic action of adrenaline and atropine methyl-nitrate in BROVON inhalant ensures speedy relief of asthma. Accurate dosage and deep inhalation are assured when used with any of our inhalers (e.g., Brovon, Deedon, Bon-Accord and Midget inhalers). This combined treatment is particularly valuable for treatment of paroxysms and for rapid relief of bronchiolar spasm often present in chronic bronchitis and emphysema.

Particulars from our Agents: **POWLEY & COMPANY (PTY.) LTD.**,
 21-24 Queens House, 11 Queen Street, Durban
 P.O. Box 4259 Cape Town P.O. Box 9628 Johannesburg
FEDERATION OF RHODESIA & NYASALAND. Agents: ASHTON & McDONALD (PVT) LTD. P.O. Box 379, Salisbury, S.R.



MOORE MEDICINAL PRODUCTS LTD
ABERDEEN LONDON OFFICE: 64 GLOUCESTER PLACE, W.1. **LONDON**

*The finest cigarette
 money can buy*



**PLAIN or with the
 MIRACLE FILTER**

Bermide

SUCCINATE-SALICYLATE THERAPY FOR THE RELIEF OF SYMPTOMS ASSOCIATED WITH ALL RHEUMATIC DISORDERS

BERMIDE fills a long-sought objective in the therapy of arthritis and rheumatic disorders. It combines the following advantages:

1 PROMPT RELIEF OF SYMPTOMS

2 PHYSIOLOGICAL ACTION

**3 FREEDOM FROM ILL-EFFECT.
OBVIATION OF SALICYLATE TOXICITY**

**4 SUITABILITY FOR PROTRACTED
ADMINISTRATION**

5 LOW COST

It has long been accepted by the medical profession that acetylsalicylic acid is unsurpassed as an adjuvant in the treatment of arthritic and rheumatic conditions.

Hitherto, it has been impossible to administer massive and prolonged dosage of acetylsalicylic acid without lowering prothrombin level and avoiding its inhibitory effect on tissue respiration.

When, however, calcium succinate and acetylsalicylic acid are combined as in BERMIDE, the acetylsalicylic acid is rendered non-toxic without lessening its effectiveness as a means of alleviating pain.


The BERMIDE formula has been enthusiastically accepted in

Great Britain, the United States and Canada, where clinical investigation has yielded impressive results.

BERMIDE is promoted ethically in bottles of 100 tablets and the large size dispensing bottle of 500 tablets. If your Pharmacist does not already have BERMIDE in stock he may obtain it directly from B. P. Davis Limited, P.O. Box 3371, Johannesburg, or through usual wholesale channels.

Bermide

BER



- Fast gaining popularity among the Medical Profession for greater convenience in operation — Quick change over from lead to lead — Continuous automatic time check — Compact and portable metal housing — Fully Guaranteed — Very reasonably priced. Backed by efficient technical service.

For further information and detailed literature write or phone:-



Union Medical Supplies LIMITED

P.O. Box 3907
Cape Town

P.O. Box 5992
Johannesburg

P.O. Box 2238
Salisbury

MEDICAL PUBLICATIONS



JUTA & CO. LTD.

Cape Town & Johannesburg

- Prof. C. F. M. Saint: An Introduction to Clinical Surgery, 3rd. Ed. .. 45s.*
Dr. M. Gelfand: Schistosomiasis .. 25s.
Ellis's Anatomy: Revised Edition by Dr. J. A. Keen .. 38s.
Dr. Louis Franklin Freed: The Problem of European Prostitution in Johannesburg .. 40s.
Marga Wright: Hygiene in South Africa 2nd. Ed. .. 17s. 6d.
A. S. Robertson: Hygiene and Sanitation in South Africa .. 30s.

The Journal of Forensic Medicine

(Sponsored by the South African Medico-Legal Society)

Published Quarterly · Annual Subscription: 42s.

A new development in antipyretic and analgesic therapy

Delton

COMPOUND

SALICYLAMIDE TABLETS

A compound tablet of the antipyretic-analgesic type, based on Salicylamide in balanced combination with Phenacetin, Codeine Phosphate and Caffeine. Each Delton tablet contains:

SALICYLAMIDE	54.16%
PHENACETIN	8.33%
CODEINE PHOSPHATE	0.91%
CAFFEINE	8.33%

INDICATIONS

DELTON can be administered in all cases in which salicylate therapy is indicated. These include acute rheumatic fever, rheumatoid arthritis, fibrositis and polyarthritis. The analgesic effect of DELTON is also of value in cases of sciatica, neuritis, myalgia, lumbago, pleurisy and neuralgia. (Reports by Stempel, German Therapeutic Congress (Karlsruhe) 1949, Wegmann T. Schweiz Med. Wschr. 1950, 80, 62. Reipert R. J. Pharmacol 1951, 76, 1312.)

DOSAGE

HEADACHES, NEURALGIA, TOOTHACHE
PERIODIC PAIN.—One tablet. Repeat 2—3 times a day, if needed.

FIBROSITIS, SCIATICA, LUMBAGO, NEURITIS
or MUSCULAR RHEUMATISM.—Two tablets. Repeat 1—2 times a day, if needed.

INFLUENZA, FEVERISH COLDS.—Two tablets, taken with a hot drink, before going to bed.

ADMINISTRATION

DELTON Tablets can be taken whole, or dispersed in water or milk. *They should not be given to children under five.* For older children, half to one tablet, according to age.

PACKING

DELTON is presented in 11.8 gr. tablets and packed in glass phials of 10 tablets, protected by a carton.

A product of

HARROGATE PHARMACEUTICALS LTD.
 Bridgend, Glam., Great Britain

Samples and further information available from

PROTEA PHARMACEUTICALS LTD.
 7 NEWTON STREET, WEMMER JOHANNESBURG
 P.O. Box 7793 Tel. 33-2211


Also at Cape Town, Port Elizabeth, East London and Durban



WATSON "MOBILIX"

WARD OUTFIT

Available now with alternative outputs



The new tube trunnion affords a full range of tube angulation.

This well-known Mobilix Outfit in its latest improved form, has the added advantage of a tube-clamp which permits the X-ray beam to be aimed in a forward direction and thus facilitates the taking of lateral radiographs of bedridden patients, views of the hip-joint in the operating theatre, etc.

For improved results in chest radiography and shorter exposures generally, the Mobilix 100, equipped with a Dynamax 20 rotating anode tube, and an output of 100 mA. is recommended.

Please ask for a copy of the second edition of Publication No. 117.

Represented in South Africa and the Rhodesias by

THE BRITISH GENERAL ELECTRIC CO. (PTY) LTD.

Box 1327, Cape Town
Box 922, Durban

Box 2406, Johannesburg

Box 914, Bloemfontein
Box 42, Port Elizabeth

THE BRITISH GENERAL ELECTRIC CO. OF CENTRAL AFRICA LTD.

Box 1070, Bulawayo

Representing

Box 845, Salisbury

THE GENERAL ELECTRIC CO. LTD. OF ENGLAND



OVER WORKED
NERVOUS
FEARFUL
TENSE
WORRIED
RESTLESS
ANXIOUS

a New Experience in **TRANQUILLITY**

For anxious, tense, restless patients Seconesin provides a safe relaxant-sedative. It introduces a totally new idea in sedation. Non-narcotic and with a minimum of secobarbital, there is no danger of cumulation or "hangover" because both components are rapidly eliminated. Seconesin acts promptly and its effect lasts for a few hours only. Day-time relaxation with Seconesin is so calming that most patients sleep well at night without further hypnotics or sedatives. Each tablet contains :—Mephenesin 400 mg. Secobarbital 30 mg.

SECONESIN

TRADE MARK

Packings: Bottles of 25, 100 and 500 tablets. Full literature on request to P.O. Box 1573, Johannesburg.

THE CROOKES LABORATORIES LIMITED • PARK ROYAL • LONDON N.W.10



Known and trusted for a hundred years



Branches at:

CAPE TOWN
DURBAN
BLOEMFONTEIN
PORT ELIZABETH
JOHANNESBURG
MOSEL BAY
EAST LONDON
KIMBERLEY

Lennon Limited, manufacturers of National Health Products, have enjoyed the confidence of the Medical Profession for over a hundred years. National Health Products have always been made to conform to the most exacting requirements of modern medicine and hygiene.

N.H.P. Products include Infalose, the well known baby food, and a range of ethical remedies and first-aid requirements for the medical profession.

LENNON • LIMITED

Chemists to South Africa

9175-1 med.



TOMORROW'S IMPROVEMENTS

TODAY

THE NEW IMPROVED BOYLE APPARATUS

MODEL 'H'
INCLUDING ABSORBER

The Boyle range of apparatus is supplied with the 'rotameter' gas measuring unit. The latest Boyle Mark II Circle Absorber can be readily fitted if closed circuit anaesthesia is required.

Latest features include: 1. 'All metal' vaporisers.

2. The new 'Trilene Interlock Unit' which is a precaution against trichlorethylene getting into the closed circuit.

(Note: All units are inter-changeable and can be bought as separate parts.)


Further details concerning this and all other medical apparatus supplied by Messrs. African Oxygen & Acetylene (Pty.) Ltd., will be gladly given on request.



AFRICAN OXYGEN & ACETYLENE (PTY) LTD MEDICAL DIVISION

76, King George Street, Hillbrow, Phone 44-4986, Johannesburg
and throughout the Union, the Rhodesias, East Africa and South West Africa

NEW
'Drinamyl Spansule'* CAPSULES
for the harassed patient



Available for prescription in containers of 30 capsules

Each STRENGTH NO. 1 capsule contains :
 10mg. 'Dexedrine' (dextro-amphetamine sulphate)
 65mg. (gr. 1) amylobarbitone

Each STRENGTH NO. 2 capsule contains :
 15mg. 'Dexedrine' (dextro-amphetamine sulphate)
 97mg. (gr. 1½) amylobarbitone

Samples and literature on request

A NEW SKF PRODUCT



SMITH KLINE & FRENCH

represented by M. & J. Pharmaceuticals (Pty.) Ltd., Diesel Street, Port Elizabeth

SUOLP1735A

'Drinamyl', 'Dexedrine' and 'Spansule' are registered trade marks

**S.A. Pat. No. 17,710*

ANNOUNCING

Nūtinal

the **new** oral
tranquillizing agent

NŪTINAL is of great value in the treatment of a variety of psychoneurotic states, especially those associated with anxiety, depression and obsessive-compulsive traits. It is particularly useful in those anxiety states which are characterized by psychosomatic symptoms such as palpitation, sweating and facial flushing.

NŪTINAL

Bottles of 100 tablets
each containing 1 mg. benactyzine hydrochloride



Made by:

BOOTS PURE DRUG COMPANY LIMITED, NOTTINGHAM, ENGLAND

and distributed by

B.P.D. (S.A.) (PTY.) LIMITED, P.O. BOX 45, JEPPESTOWN, TRANSVAAL

BPD/NU/16

7 July 1956

MEDICAL PROCEEDINGS • MEDIESE BYDRAES

XXXV

polycycline

aqueous pediatric drops

MAXIMUM TOLERANCE pH 7.5 (NEUTRAL)

READY-TO-USE • STABLE

Miscible with orange juice, formula or milk.

Stable for 24 months at room temperature.

Calibrated dropper supplied with 10 cc. bottle for accurate dosage and easy administration.

INDICATED against a large variety of Gram-positive and Gram-negative organisms, certain rickettsiae, large viruses and organisms resistant to other antibiotics.

DOSAGE is approximately 10 mg. per pound of body weight per 24 hours.

POLYCYCLINE is also available in the following forms: Capsules, Suspension with Triple-Sulphonamides, Dermatologic and Ophthalmic Ointments, Intra-muscular & Intravenous Injections.


Samples and Literature available from:

BRISTOLABS (PTY.) LTD.
P.O. Box 2515
JOHANNESBURG



Bristol
LABORATORIES INC.
SYRACUSE, N.Y.

Depot Stockists: Heynes Mathew, Limited, Cape Town; South African Drug Houses, Durban; E. P. Drug House, Port Elizabeth; James Reid (Pty.) Limited, Bloemfontein; Cloete Kruger (Pty.) Limited, Windhoek.



When
appetite fails
in old age and
convalescence

ELIXIR
Gerix

ABBOTT'S GERIATRIC ELIXIR

Pleasantly wine-flavoured Elixir GERIX provides effective therapy when appetites fall behind the continuing need for nutrition. Taken before meals, GERIX acts as a stomachic, increasing the desire for food. At the same time, GERIX provides important nutritional elements—B-complex factors and iron—often lacking in faulty diets.

New
pleasant
tasting
appetite stimulant
and nutritional
supplement



Abbott


JOHANNESBURG: 223-5, Booysens Road, Selby. (P.O. Box 1616) Telephone 33-4556 7/8; DURBAN, NATAL: Maiglen House, First Floor, 71 Gale Street (P.O. Box 2291) Telephone 68089; BLOEMFONTEIN: 2, Short Street, (P.O. Box 731) O.F.S. Telephone 81457; CAPE TOWN: Marine Chambers, 4, Lower Burg Street (P.O. Box 2923) Telephone 3-0659; PRETORIA: Colorado, 115 Schoeman Street, Telephone 38800-38805; PORT ELIZABETH: 30 Crawford Street (P.O. Box 3186) Tele. 87610
EAST LONDON: 115 Oxford Street, (P.O. Box 994) Telephone 72806
All Telegraphic Addresses: "Abbottlab"



WATSON "MOBILIX"

WARD OUTFIT

Available now with alternative outputs



The new tube trunnion affords a full range of tube angulation.

This well-known Mobilix Outfit in its latest improved form, has the added advantage of a tube-clamp which permits the X-ray beam to be aimed in a forward direction and thus facilitates the taking of lateral radiographs of bedridden patients, views of the hip-joint in the operating theatre, etc.

For improved results in chest radiography and shorter exposures generally, the Mobilix 100, equipped with a Dynamax 20 rotating anode tube, and an output of 100 mA. is recommended.

Please ask for a copy of the second edition of Publication No. 117.

Represented in South Africa and the Rhodesias by

THE BRITISH GENERAL ELECTRIC CO. (PTY) LTD.

Box 1327, Cape Town
Box 922, Durban

Box 2406, Johannesburg

Box 914, Bloemfontein
Box 42, Port Elizabeth

THE BRITISH GENERAL ELECTRIC CO. OF CENTRAL AFRICA LTD.

Box 1070, Bulawayo

Representing

Box 845, Salisbury

THE GENERAL ELECTRIC CO. LTD. OF ENGLAND



OVER WORKED
NERVOUS
TENSE
FEARFUL
WORRIED
RESTLESS
ANXIOUS

a New Experience in
TRANQUILLITY

For anxious, tense, restless patients Seconesin provides a safe relaxant-sedative. It introduces a totally new idea in sedation. Non-narcotic and with a minimum of secobarbital, there is no danger of cumulation or "hangover" because both components are rapidly eliminated. Seconesin acts promptly and its effect lasts for a few hours only. Day-time relaxation with Seconesin is so calming that most patients sleep well at night without further hypnotics or sedatives. Each tablet contains :—Mephenesin 400 mg. Secobarbital 30 mg.

SECONESIN

TRADE MARK

Packings: Bottles of 25, 100 and 500 tablets. Full literature on request to P.O. Box 1573, Johannesburg.

THE CROOKES LABORATORIES LIMITED • PARK ROYAL • LONDON N.W.10



**Known and trusted for
a hundred years**



Branches at:

CAPE TOWN
DURBAN
BLOEMFONTEIN
PORT ELIZABETH
JOHANNESBURG
MOSSEL BAY
EAST LONDON
KIMBERLEY

Lennon Limited, manufacturers of National Health Products, have enjoyed the confidence of the Medical Profession for over a hundred years. National Health Products have always been made to conform to the most exacting requirements of modern medicine and hygiene.

N.H.P. Products include Infalose, the well known baby food, and a range of ethical remedies and first-aid requirements for the medical profession.

LENNON • LIMITED

Chemists to South Africa



TOMORROW'S IMPROVEMENTS

TODAY

THE NEW IMPROVED BOYLE APPARATUS

MODEL 'H'
INCLUDING ABSORBER

The Boyle range of apparatus is supplied with the 'rotameter' gas measuring unit. The latest Boyle Mark II Circle Absorber can be readily fitted if closed circuit anaesthesia is required.

Latest features include: 1. 'All metal' vaporisers.

2. The new 'Trilene Interlock Unit' which is a precaution against trichlorethylene getting into the closed circuit.

(Note: All units are inter-changeable and can be bought as separate parts.)

Further details concerning this and all other medical apparatus supplied by Messrs. African Oxygen & Acetylene (Pty.) Ltd., will be gladly given on request.



AFRICAN OXYGEN & ACETYLENE (PTY) LTD MEDICAL DIVISION

76, King George Street, Hillbrow, Phone 44-4998, Johannesburg
and throughout the Union, the Rhodesias, East Africa and South West Africa

NEW

'Drinamyl Spansule'* CAPSULES

for the harassed patient



Available for prescription in containers of 30 capsules

<p>Each STRENGTH NO. 1 capsule contains :</p> <p>10mg. 'Dexedrine' (dextro-amphetamine sulphate)</p> <p>65mg. (gr. 1) amylobarbitone</p>	<p>Each STRENGTH NO. 2 capsule contains :</p> <p>15mg. 'Dexedrine' (dextro-amphetamine sulphate)</p> <p>97mg. (gr. 1½) amylobarbitone</p>
---	--

Samples and literature on request

A NEW SKF PRODUCT



SMITH KLINE & FRENCH

represented by M. & J. Pharmaceuticals (Pty.) Ltd., Diesel Street, Port Elizabeth

SUDLP1158A

'Drinamyl', 'Dexedrine' and 'Spansule' are registered trade marks

*S.A. Pat. No. 17,710

ANNOUNCING

Nūtinal

the **new** oral
tranquillizing agent

NŪTINAL is of great value in the treatment of a variety of psychoneurotic states, especially those associated with anxiety, depression and obsessive-compulsive traits. It is particularly useful in those anxiety states which are characterized by psychosomatic symptoms such as palpitation, sweating and facial flushing.

NŪTINAL

Bottles of 100 tablets
each containing 1 mg. benactyzine hydrochloride



Made by:

BOOTS PURE DRUG COMPANY LIMITED, NOTTINGHAM, ENGLAND

and distributed by

B.P.D. (S.A.) (PTY.) LIMITED, P.O. BOX 45, JEPPESTOWN, TRANSVAAL

BPD/NUJ/16

polycycline

aqueous pediatric drops

MAXIMUM TOLERANCE pH 7.5 (NEUTRAL)

READY-TO-USE • STABLE

Miscible with orange juice, formula or milk.

Stable for 24 months at room temperature.

Calibrated dropper supplied with 10 cc. bottle for accurate dosage and easy administration.

INDICATED against a large variety of Gram-positive and Gram-negative organisms, certain rickettsiae, large viruses and organisms resistant to other antibiotics.

DOSAGE is approximately 10 mg. per pound of body weight per 24 hours.

POLYCYCLINE is also available in the following forms: Capsules, Suspension with Triple-Sulphonamides, Dermatologic and Ophthalmic Ointments, Intra-muscular & Intravenous Injections.


Samples and Literature available from:

BRISTOLABS (PTY.) LTD.
P.O. Box 2515
JOHANNESBURG



Bristol
LABORATORIES INC.
SYRACUSE, N.Y.

Depot Stockists: Heynes Mathew, Limited, Cape Town; South African Drug Houses, Durban; E. P. Drug House, Port Elizabeth; James Reid (Pty.) Limited, Bloemfontein; Cloete Kruger (Pty.) Limited, Windhoek.




When
appetite fails
in old age and
convalescence



ABBOTTS GERIATRIC ELIXIR

Pleasantly wine-flavoured Elixir GERIX provides effective therapy when appetites fall behind the continuing need for nutrition. Taken before meals, GERIX acts as a stomachic, increasing the desire for food. At the same time, GERIX provides important nutritional elements—B-complex factors and iron—often lacking in faulty diets.

New
pleasant
tasting
appetite stimulant
and nutritional
supplement



JOHANNESBURG: 223-5, Booysens Road, Selby. (P.O. Box 1816) Telephone 33-4556 7/8; **DURBAN, NATAL:** Maiglen House, First Floor, 71 Gale Street (P.O. Box 2291) Telephone 68088; **BLOEMFONTEIN:** 2, Short Street, (P.O. Box 731) O.F.S. Telephone 81457; **CAPE TOWN:** Marine Chambers, 4, Lower Burg Street (P.O. Box 2923) Telephone 3-0559; **PRETORIA:** Colorado, 115 Schoeman Street, Telephone 38900—38905; **PORT ELIZABETH:** 30 Crawford Street (P.O. Box 3186) Tele. 87810
EAST LONDON: 118 Oxford Street, (P.O. Box 984) Telephone 72805
All Telegraphic Addresses: "Abbottlab"